

The Chern Character of ϑ -summable Fredholm Modules over dg Algebras and Localization on Loop Space

Batu Güneysu* and Matthias Ludewig†

July 25, 2019

Abstract

We introduce the notion of a ϑ -summable Fredholm module over a locally convex dg algebra Ω and construct its Chern character as a cocycle on the entire cyclic complex of Ω , extending the construction of Jaffe, Lesniewski and Osterwalder to a differential graded setting. Using this Chern character, we prove an index theorem involving an abstract version of a Bismut-Chern character constructed by Getzler, Jones and Petrack in the context of loop spaces. Our theory leads to a rigorous construction of the path integral for $N = 1/2$ supersymmetry which satisfies a Duistermaat-Heckman type localization formula on loop space, solving a long standing open problem.

Contents

1	Introduction	2
2	ϑ-summable Fredholm Modules over Locally Convex dg Algebras	7
3	Algebraic Preliminaries	11
3.1	Bar Chains and Cochains	11
3.2	Cyclic Chains and Cochains	12
3.3	Analytic Cochains and Entire Chains	13
3.4	Chen Normalization	15
4	The Quantization Map	16
5	The Chern Character	24

*Universität Bonn. E-mail: guneysu@math.uni-bonn.de

†The University of Adelaide. E-mail: matthias.ludewig@adelaide.edu.au

6	Homotopy Invariance of the Chern Character	28
7	The Bismut-Chern Character of an Idempotent	31
8	An Index Theorem	35
9	The Supersymmetric Path Integral and its Localization Property	38

1 Introduction

The close relation between supersymmetry and index theory was discovered by Alvarez-Gaumé [1, 2], motivated by considerations of Witten [52, 53], and has gotten a lot of attention ever since (see e.g. [3, 5, 10, 11, 27, 26, 28, 34, 31, 30, 39, 43, 49]). In particular, Alvarez-Gaumé observed that the path integral for the $N = 1/2$ supersymmetric σ -model enables a concise proof of the Atiyah-Singer index theorem for the Dirac operator on a spin manifold, by *bona fide* generalizing a the Duistermaat-Heckman localization formula to the (infinite-dimensional) manifold of smooth loops. Much more than obtaining another proof of the Atiyah-Singer index theorem, these considerations suggest to study the following fundamental question:

*To what extent does one have a localization formula
on the loop space of a spin manifold?*

Note that most of the above mentioned calculations had to remain formal so far, due to the mathematically ill-defined nature of the supersymmetric path integral: Even though the integral in question has a Gaussian nature (cf. formula (1.1) below), the usual approach (which uses continuous loops instead of smooth loops and the Wiener measure for the bosonic integration) makes it notoriously hard to implement supersymmetry.

This article provides a framework that for the first time allows to construct the above path integral rigorously for a large class of integrands, in such a way that it is supersymmetric and the localization formula holds. This finally justifies the above formal calculations after almost 35 years.

To set the stage, let X be compact, even-dimensional spin manifold X . In the differential geometric formulation of Alvarez-Gaumé's observations by Atiyah [5] and Bismut [10], the path integral of interest is the integration functional

$$I[\xi] \stackrel{\text{formally}}{=} \int_{\text{LX}} e^{-S+\omega} \wedge \xi \tag{1.1}$$

on smooth differential forms ξ on the space LX of smooth loops in X , where S is the energy functional and ω is the canonic presymplectic form on LX . The *supersymmetry* of the path integral is then the formula $I[(d + \iota_K)\xi] = 0$, where ι_K denotes the contraction by the velocity vector field $K(\gamma) = \dot{\gamma}$ which generates the natural S^1 -action on LX given by rotating each loop.

While the above expression for I is not mathematically well-defined, one expects to have the *localization formula*¹

$$I[\xi] = (2\pi)^{-\dim(X)/2} \int_X \widehat{A}(X) \wedge \xi|_X \quad (1.2)$$

for differential forms ξ that are equivariantly closed, $(d + \iota_K)\xi = 0$, in analogy with the results of Duistermaat-Heckman and Berline-Vergne [25, 8] on finite-dimensional integrals of this kind (cf. the arguments of Atiyah in [5]). At this point, the proof of the Atiyah-Singer index theorem for twisted Dirac operators becomes a simple consequence of supersymmetry and localization, using the special integrands introduced by Bismut [10].

Our approach to define such an integration functional I is based on Chen's iterated integral map [18, 19, 20]

$$\rho : \mathbb{C}(\Omega(X)) \longrightarrow \Omega(LX)$$

and its extensions (c.f. [34, 17] and §9), which constructs differential forms on LX from elements of the *cyclic chain complex* $\mathbb{C}(\Omega(X))$ of the differential graded algebra $\Omega(X)$ (c.f. §3). Hence instead of working on $\Omega(LX)$, we construct a closed functional on $\mathbb{C}(\Omega(X))$, which can then be pushed forward using the iterated integral map to obtain an integration functional I , defined on all differential forms on LX that are iterated integrals². The supersymmetry of this path integral then follow from the closedness of the cochain; moreover, we provide a complete proof of the loop space localization formula (1.2).

One of the main new insights of this paper is that the cochain on $\mathbb{C}(\Omega(X))$ used to define the path integral can be seen as a *Chern character*, defined in terms of the heat semigroup of the Dirac operator on X (c.f. Example 2.3) below). This dependence reflects the fact that the orientability of LX is connected to the spin property of X [48, 50].

Moreover, it turns out that in the construction of this Chern character, $\Omega(X)$ may be replaced by a general (not necessarily graded commutative) locally convex dg algebra Ω . In fact, it turns out that even for the definition of I , it is useful to have the general theory at disposal (see e.g. the proof of Thm. 8.1, related to Example 2.2). The Chern character is then associated to a structure over Ω , which we call a *ϑ -summable Fredholm module* over Ω . Surprisingly, our Chern character generalizes that of Jaffe, Lesniewski and Osterwalder [39] in the ungraded case³

When Ω is a unital (ungraded) Banach algebra, such a Fredholm module (as introduced by Connes in [21, 22]) consists of a representation \mathfrak{c} of Ω on a \mathbb{Z}_2 -graded Hilbert space \mathcal{H} together with an odd self-adjoint unbounded operator Q on \mathcal{H} (satisfying several analytic conditions). When generalizing the notion of a Fredholm module to this setting, it turns out that requiring \mathfrak{c} to be a representation of Ω turns out to be too restrictive for the applications we have in mind. Instead, we just assume that \mathfrak{c} is a degree-preserving linear

¹This in fact implies the supersymmetry property $I[(d + \iota_K)\xi] = 0$.

²The restriction to forms having a certain integrability property seems only natural, as even on finite dimensional *noncompact* manifolds one cannot integrate every smooth differential form.

³In the sense that if $\Omega = \Omega^0$ is concentrated in degree zero, then $\text{Ch}(\mathcal{M})$ is equal to the JLO cocycle.

map from Ω to $\mathcal{L}(\mathcal{H})$ such that⁴

$$[Q, \mathbf{c}(f)] = \mathbf{c}(df), \quad \text{and} \quad \mathbf{c}(f\theta) = \mathbf{c}(f)\mathbf{c}(\theta), \quad \mathbf{c}(\theta f) = \mathbf{c}(\theta)\mathbf{c}(f). \quad (1.3)$$

for all $f \in \Omega^0$ and all $\theta \in \Omega$; in particular, \mathbf{c} is required to be a representation only on the subalgebra Ω^0 . These assumptions are satisfied by the Fredholm module \mathcal{M}^X over X , where \mathbf{c} stems from Clifford multiplication and $Q = \mathbf{D}$ is the Dirac operator.

Given such an abstract Fredholm module \mathcal{M} , its *Chern character* is then a linear functional $\text{Ch}(\mathcal{M})$ on the *cyclic chain complex* $\mathbf{C}(\Omega)$ of Connes. This complex carries three differentials: the differential \underline{d} , which is induced from the differential d of Ω , the *Hochschild differential* \underline{b} and the *Connes operator* \underline{B} . The Chern character is closed with respect to the *total differential* $\underline{d} + \underline{b} + \underline{B}$ (c.f. Thm. 5.3 below) which acts on linear functionals on $\mathbf{C}(\Omega)$ by precomposition, c.f. (1.7).

Theorem A. *The Chern character $\text{Ch}(\mathcal{M})$ is even and closed, that is,*

$$(\underline{d} + \underline{b} + \underline{B})\text{Ch}(\mathcal{M}) = 0. \quad (1.4)$$

Let us briefly outline our construction of the Chern character, in a presentation similar to that of Quillen [47, §9] in the ungraded case (c.f. also [46]). Following Quillen, an odd bar cochain ω on Ω (c.f. (3.1) below) with values in an algebra L can be seen as a *connection form* on the space of L -valued bar cochains. Its *curvature* is then given by

$$F = \delta\omega + \omega^2,$$

where δ is the codifferential on $\mathbf{B}(\Omega)$. The observation is now that a ϑ -summable Fredholm module \mathcal{M} over Ω determines such a cochain $\omega = \omega_{\mathcal{M}}$ taking values in linear operators on \mathcal{H} , with curvature $F_{\mathcal{M}}$ (c.f. (4.3) and (4.4) below); the Chern character is then defined by

$$\langle \text{Ch}(\mathcal{M}), (\theta_0, \dots, \theta_N) \rangle := \text{Str} \left(\mathbf{c}(\theta_0) \Phi^{\mathcal{M}}(\theta_1, \dots, \theta_N) \right), \quad \text{with} \quad \Phi^{\mathcal{M}} = \exp(-F_{\mathcal{M}}),$$

in spirit of Chern-Weil theory. Closedness of $\text{Ch}(\mathcal{M})$, Thm. A, is then essentially a consequence of the Bianchi identity for $F_{\mathcal{M}}$. However, the fact that $\omega_{\mathcal{M}}$ and $F_{\mathcal{M}}$ take values in *unbounded* operators, not in $\mathcal{L}(\mathcal{H})$, poses severe analytical problems; tackling these belongs to the main tasks of this paper.

Related to these analytic issues is the important feature of our Chern character that its homogeneous components decay rapidly enough to extend it to a continuous linear functional on the *entire complex* $\mathbf{C}^e(\Omega)$, which contains infinite sums of homogeneous chains, subject to certain growth conditions modelled on the growth of entire functions (c.f. Thm. 5.2 below). We remark that this growth condition was first introduced on *cochains* by Connes in [21]; in our case, it appears naturally on *chains*, as in [35]. Motivated by the terminology of Meyer [44], we call the dual growth condition satisfied by $\text{Ch}(\mathcal{M})$ *analytic*, as it is related to the growth of analytic functions (cf. §3.3).

⁴In fact, not even these algebraic properties are needed; dropping these assumptions leads to what we call a *weak* Fredholm module, c.f. §2.

Theorem B. *The Chern character $\text{Ch}(\mathcal{M})$ is analytic, that is, it extends to a continuous linear functional on the entire complex $\mathbb{C}^\epsilon(\Omega)$.*

To prove an index theorem for our Chern character, we use the Bismut-Chern character $\text{Ch}(p)$ constructed by Getzler, Jones and Petrack in [34]. It is an entire chain, which allows to pair it with our Chern character in view of Thm. B. An obstacle is that $\text{Ch}(p)$ is not a chain over Ω but over its *acyclic extension*

$$\Omega_{\mathbb{T}} := \Omega[\sigma],$$

where σ is a formal variable of degree -1 satisfying $\sigma^2 = 0$. Now it turns out that any ϑ -summable Fredholm module \mathcal{M} over Ω can be extended to a ϑ -summable weak Fredholm module $\mathcal{M}_{\mathbb{T}}$ over $\Omega_{\mathbb{T}}$, giving rise to a Chern character $\text{Ch}(\mathcal{M}_{\mathbb{T}})$ (c.f. Example 2.5 below). The following result (c.f. Thm. 8.1 below) then generalizes the index theorem of Getzler and Szenes [35, Thm. D].

Theorem C. *For every idempotent $p \in \text{Mat}_n(\Omega^0)$ we have the index formula*

$$\text{ind}(Q_p) = \langle \text{Ch}(\mathcal{M}_{\mathbb{T}}), \text{Ch}(p) \rangle. \quad (1.5)$$

While $\text{Ch}(p)$ is not closed in the complex $\mathbb{C}^\epsilon(\Omega_{\mathbb{T}})$, it is closed in the quotient complex

$$\mathbf{N}^{\mathbb{T},\epsilon}(\Omega) = \mathbb{C}^\epsilon(\Omega_{\mathbb{T}}) / \overline{\mathbf{D}^{\mathbb{T}}(\Omega)}$$

of *Chen normalized chains*, which is central to our theory (c.f. Def. 3.4 below). It turns out that the properties (1.3) ensure that $\text{Ch}(\mathcal{M}_{\mathbb{T}})$ is *Chen normalized*, meaning that it vanishes on $\mathbf{D}^{\mathbb{T}}(\Omega)$ and therefore descends to a functional on $\mathbf{N}^{\mathbb{T},\epsilon}(\Omega)$ (Thm. 5.5 below). This leads to a cohomological version of the index theorem involving the K -theory of the algebra Ω^0 (Corollary 8.3).

We emphasize that the Chern-Weil type method outlined above constructs the Chern character as a *cochain*, not only as a cohomology class. A remarkable feature of this cochain is that the pairing on the right hand side of (1.5) is *exactly equal* to the supertrace of the heat operator $e^{-Q_p^2}$ (c.f. Prop. 8.1); using this, the proof of Thm. C does not need the homotopy invariance of $\text{Ch}(\mathcal{M}_{\mathbb{T}})$, and follows straightforwardly from the McKean-Singer formula. Nevertheless, we prove:

Theorem D. *The Chern character $\text{Ch}(\mathcal{M}_{\mathbb{T}})$ is invariant under homotopies of \mathcal{M} .*

In fact, the Chern character is even homotopy-invariant as a *Chen normalized cochain*, i.e. as one on the complex $\mathbf{N}^{\mathbb{T},\epsilon}(\Omega)$; for a more precise statement, see §6, where this result is obtained from a Chern-Simons type transgression formula.

Making contact with the construction of the functional I , taking $\Omega = \Omega(X)$ and $\mathcal{M} = \mathcal{M}^X$, the Fredholm module over $\Omega(X)$ determined by the Dirac operator (c.f. Example 2.3) it follows that we can define the path integral map by the formula

$$I[\rho(c)] = i^{\dim(X)/2} \langle \text{Ch}(\mathcal{M}_{\mathbb{T}}^X), c \rangle \quad (1.6)$$

for all $c \in \mathcal{C}(\Omega(X)_{\mathbb{T}})$, noting that $\mathcal{C}(\Omega(X)_{\mathbb{T}})$ is the natural domain of definition of the extended version of Chen's iterated integral map. Given the basic properties [17] of ρ , the well-definedness, respectively supersymmetry of I turns out to follow from the fact that $\text{Ch}(\mathcal{M}_{\mathbb{T}}^X)$ is Chen normalized, respectively closed, Thm. A. The localization formula (1.2) for the path integral is a consequence of the following result, which is proved using Thm. D in combination with a version of Getzler-rescaling.

Theorem E. *For a compact, even-dimensional spin manifold X , let \mathcal{M}^X be the Fredholm module over $\Omega(X)$ given by the Dirac operator. Then as an element of the complex $\mathbb{N}^{\mathbb{T}, \epsilon}(\Omega(X))$, the Chern character $\text{Ch}(\mathcal{M}_{\mathbb{T}}^X)$ is cohomologous to the cochain μ_0 given by the formula*

$$\mu_0(\theta_0, \dots, \theta_N) = \frac{1}{(2\pi i)^{\dim(X)/2} N!} \int_X \widehat{A}(X) \wedge \theta'_0 \wedge \theta''_1 \wedge \dots \wedge \theta''_N,$$

for all $\theta_j = \theta'_j + \sigma\theta''_j \in \Omega(X)_{\mathbb{T}}$, where $\widehat{A}(X)$ is the \widehat{A} -form given in (9.2).

The analyticity of $\text{Ch}(\mathcal{M}_{\mathbb{T}}^X)$, Thm. B, allows to integrate the Bismut-Chern characters introduced by Bismut in [10], which together with Thm. C and the localization formula enables a rigorous version of the proof of the twisted Atiyah-Singer theorem envisioned in [4, 10].

These results show that our integral map I has the desired properties on a cohomological level. Geometrically, one can ask for the relation of the map I defined by (1.6) to the heuristic formula (1.1). This question is answered in [36, 37], where an integral map is constructed directly on $\Omega(LX)$, by a suitable interpretation of (1.6) in terms of regularized Pfaffians and the Wiener measure. The result is then that this map is *exactly equal* to the path integral map constructed in this article. We expect that this other description permits a probabilistic proof of the localization principle along the lines of [9, 6].

To the best of our knowledge, the idea to use cyclic cohomology of Ω and the iterated integral map to construct the supersymmetric path integral is due to Getzler, and this paper can be seen as a completion of this endeavor. Needless to say, our paper is largely inspired by his work (in particular [35, 34, 31, 30, 32]). Moreover, Bismut's results suggest a possible connection between our machinery and the hypoelliptic Laplacian, respectively the hypoelliptic Dirac operator [12, 13, 14].

We would also like to remark that our results could also be stated in the abstract framework of bornological algebras instead of the locally convex setup, as in the work of Meyer [44]. However, we believe that the use of locally convex algebras somewhat simplifies our presentation; in particular the entire growth condition on chains takes a somewhat technical form in the bornological setup (cf. §3.3). Moreover, we observe that our main example $\Omega = \Omega(X)$, X a compact manifold, is a nuclear Fréchet algebra, so that the von-Neumann and the precompact bornology coincide.

The outline of this paper is as follows. First we give the definition of a ϑ -summable Fredholm module over Ω and give several motivating examples. In §3, we introduce the algebraic preliminaries for the constructions in §4, 5 and 6, where we prove Thm. A,

Thm. B and Thm. D. Then in §7, we discuss the Bismut-Chern character of an idempotent and subsequently prove Thm. C, in §8. Finally, in §9, we discuss the construction of the supersymmetric path integral. In particular, we give a proof of the localization formula (1.2), c.f. Thm. 9.1 and Thm. 9.5.

Acknowledgements. It is our pleasure to thank J.-M. Bismut, S. Cacciatori, F. Hanisch, N. Higson, M. Lesch, V. Mathai and S. Shen for helpful discussions. We are further indebted to the Max-Planck-Institute in Bonn, the Humboldt University in Berlin and The University of Adelaide, where parts of the work on this project was conducted. The second-named author was supported by the Max-Planck-Foundation and the ARC Discovery Project grant FL170100020 under Chief Investigator and Australian Laureate Fellow Mathai Varghese.

Conventions. In this paper, all vector spaces and algebras are over \mathbb{C} , and we always work in the category of *graded* vector spaces. Every \mathbb{Z} -graded vector space is considered as \mathbb{Z}_2 -graded in terms of the induced even/odd grading, and tensor products and direct sums of graded vector spaces are equipped with their canonically given \mathbb{Z} -grading (and thus the induced \mathbb{Z}_2 -grading). An ungraded vector space will be considered to be \mathbb{Z} -graded (and thus \mathbb{Z}_2 graded) by declaring all its elements to have degree zero.

If V, W are \mathbb{Z}_2 -graded vector spaces, the vector space $\text{Hom}(V, W)$ is \mathbb{Z}_2 -graded in the usual way. Given coefficients W , any $A \in \text{End}(V)$ has a \mathbb{Z}_2 -graded dual map $A^\vee \in \text{Hom}(V, W)$ given by

$$(A^\vee \ell)(v) := (-1)^{|A||\ell|} \ell(A(v)), \quad \ell \in \text{Hom}(V, W), \quad v \in V. \quad (1.7)$$

We always use this dual (instead of the ungraded version), and usually, we just write A again instead of A^\vee . Throughout, given $A, B \in \text{End}(V)$, the bracket $[A, B] \in \text{End}(V)$ denotes the graded commutator (or supercommutator)

$$[A, B] := AB - (-1)^{|A||B|} BA.$$

There are no ungraded commutators in this paper.

2 ϑ -summable Fredholm Modules over Locally Convex dg Algebras

Recall that a differential graded algebra Ω (or dg algebra for short) is an algebra that is the direct sum of subspaces $\Omega^j \subset \Omega$, $j \in \mathbb{Z}$, such that $\Omega^i \Omega^j \subset \Omega^{i+j}$ for all $i, j \in \mathbb{Z}$, together with a degree +1 differential d which satisfies the graded Leibniz rule. We always assume that Ω has a unit $\mathbf{1} \in \Omega$.

A *locally convex dg algebra* is a dg algebra Ω , where the underlying vector space carries the structure of a locally convex vector space such that the differential is continuous and such that the product map is jointly continuous. Explicitly, these two conditions mean

that for each continuous seminorm ν on Ω , there exists a continuous seminorm ν' on Ω such that

$$\nu(d\theta) \leq \nu'(\theta), \quad \nu(\theta_1\theta_2) \leq \nu'(\theta_1)\nu'(\theta_2) \quad \text{for all } \theta, \theta_1, \theta_2 \in \Omega. \quad (2.1)$$

Moreover, we require that Ω is the *topological* direct sum of its homogenous summands Ω^j ; in particular, each of these is a closed subspace of Ω .

Definition 2.1 (ϑ -summable Fredholm module). An (*even*) ϑ -summable Fredholm module over a locally convex dg algebra Ω is a triple $\mathcal{M} = (\mathcal{H}, \mathbf{c}, Q)$, where

- (i) \mathcal{H} is a \mathbb{Z}_2 -graded Hilbert space;
- (ii) $\mathbf{c} : \Omega \rightarrow \mathcal{L}(\mathcal{H})$ is a parity-preserving, bounded linear map;
- (iii) Q is an odd self-adjoint unbounded operator on \mathcal{H} ,

such that we have

$$[Q, \mathbf{c}(f)] = \mathbf{c}(df), \quad \text{and} \quad \mathbf{c}(f\theta) = \mathbf{c}(f)\mathbf{c}(\theta), \quad \mathbf{c}(\theta f) = \mathbf{c}(\theta)\mathbf{c}(f) \quad (2.2)$$

for all $f \in \Omega^0$ and all $\theta \in \Omega$. Moreover, we have the following analytic requirements.

(A1) For each $\theta \in \Omega$ the operators $C_{\pm}(\theta) := \Delta^{\pm 1/2}\mathbf{c}(\theta)\Delta^{\mp 1/2}$ are densely defined and bounded, where $\Delta = Q^2 + 1$; moreover, the assignment $\theta \mapsto C_{\pm}(\theta)$ is bounded from Ω to $\mathcal{L}(\mathcal{H})$;

(A2) For each $T > 0$ the operator $e^{-TQ^2} \in \mathcal{L}(\mathcal{H})$ is trace-class.

If instead of (2.2), \mathcal{M} only satisfies the weaker property $\mathbf{c}(\mathbf{1}) = 1$ (the identity operator), we call \mathcal{M} a ϑ -summable weak Fredholm module.

In the remainder of this section we give several examples of Fredholm modules. The following example highlights the relation to the usual notion of a ϑ -summable Fredholm module over an algebra.

Example 2.2 (Non-commutative differential forms). Let \mathcal{A} be an ungraded locally convex algebra, together with a ϑ -summable Fredholm module over \mathcal{A} in the sense of Connes [22]. That is, we are given a triple $\mathcal{M}_0 = (\mathcal{H}, \mathbf{c}_0, Q)$, where \mathcal{H} is a \mathbb{Z}_2 -graded Hilbert space, Q is an odd selfadjoint unbounded operator satisfying (A2) and $\mathbf{c}_0 : \mathcal{A} \rightarrow \mathcal{L}^+(\mathcal{H})$ a representation such that there exists a continuous seminorm ν on \mathcal{A} with

$$\|\mathbf{c}_0(a)\| + \|[Q, \mathbf{c}_0(a)]\| \leq \nu(a) \quad (2.3)$$

for all $a \in \mathcal{A}$. Under the additional assumption that there exists a continuous seminorm ν' on \mathcal{A} such that

$$\|\Delta^{1/2}[Q, \mathbf{c}_0(a)]\Delta^{-1/2}\| + \|\Delta^{1/2}[Q, \mathbf{c}_0(a)^*]\Delta^{-1/2}\| \leq \nu'(a), \quad (2.4)$$

where $\Delta = Q^2 + 1$, there exists a canonical extension of \mathcal{M}_0 to a Fredholm module $\mathcal{M} = (\mathcal{H}, \mathbf{c}, Q)$ over the differential graded algebra of non-commutative differential forms $\Omega_{\mathcal{A}}$, as we now explain.

Let us start by reviewing the construction of $\Omega_{\mathcal{A}}$. First let $\Omega_{\mathcal{A}}^0 := \mathcal{A}$ and let $\Omega_{\mathcal{A}}^1$ be the bimodule of *one-forms on \mathcal{A}* , defined as the quotient of the free bimodule on generators da , $a \in \mathcal{A}$, by the relations

$$\begin{aligned} d(\lambda a + \mu b) &= \lambda da + \mu db & a, b \in \mathcal{A}, \quad \lambda, \mu \in \mathbb{C}. \\ d(ab) &= a db + da b, \end{aligned}$$

Now we set

$$\Omega_{\mathcal{A}}^n := \underbrace{\Omega_{\mathcal{A}}^1 \otimes_{\mathcal{A}} \cdots \otimes_{\mathcal{A}} \Omega_{\mathcal{A}}^1}_n \quad \text{and} \quad \Omega_{\mathcal{A}} := \bigoplus_{n=0}^{\infty} \Omega_{\mathcal{A}}^n.$$

The product in $\Omega_{\mathcal{A}}$ is the obvious one, induced by the tensor algebra, while the formula

$$d(a_0 da_1 \cdots da_n) = da_0 da_1 \cdots da_n$$

gives a well-defined differential which turns $\Omega_{\mathcal{A}}$ into a dg algebra. Here we use that using the second relation above, any element $\theta \in \Omega_{\mathcal{A}}^n$ can be written as a sum of elements of the form $a_0 da_1 \cdots da_n$ for $a_0, \dots, a_n \in \mathcal{A}$ (for a reference, see e.g. [42, 2.6.4]).

Now it can be checked that setting

$$\mathbf{c}(a_0 da_1 \cdots da_n) := \mathbf{c}_0(a_0)[Q, \mathbf{c}_0(a_1)] \cdots [Q, \mathbf{c}_0(a_n)] \quad (2.5)$$

is a well-defined linear map $\Omega_{\mathcal{A}} \rightarrow \mathcal{L}(\mathcal{H})$ which is obviously grading preserving (as Q is odd) and satisfies (2.2). Clearly, the condition (A1) follows from (2.3) and (A2) follows since

$$\Delta^{1/2} \mathbf{c}(a_0 da_1 \cdots da_n) \Delta^{-1/2} = (\Delta^{1/2} \mathbf{c}_0(a_0) \Delta^{-1/2}) \prod_{k=1}^n (\Delta^{1/2} [Q, \mathbf{c}_0(a_k)] \Delta^{-1/2}),$$

which can be estimated using (2.3) respectively (2.4).

The following example is the main motivation for this paper.

Example 2.3 (Differential forms and spinors). Let X be an even-dimensional compact spin (or spin^c) manifold and let $\Sigma \rightarrow X$ be the corresponding complex spinor bundle with the Dirac operator D . There is a standard way to construct a ϑ -summable Fredholm module $\mathcal{M}^X = (\mathcal{H}, Q, \mathbf{c})$ over $\Omega(X)$, the dg algebra of differential forms on X , a nuclear Fréchet algebra, due to compactness of X . Here $\mathcal{H} = L^2(X, \Sigma)$, the space of square-integrable sections of Σ over X and $Q = D$, the Dirac operator, while \mathbf{c} is the so-called *quantization map* (c.f. [7, Prop. 3.5]): For differential one forms $\theta_1, \dots, \theta_k \in \Omega^1(X)$, the endomorphism of the spinor bundle $\mathbf{c}(\theta_1 \wedge \cdots \wedge \theta_k)$ is given by

$$\mathbf{c}(\theta_1 \wedge \cdots \wedge \theta_k) = \frac{1}{k!} \sum_{\sigma \in S_k} \mathbf{c}_1(\theta_{\sigma_1}) \cdots \mathbf{c}_1(\theta_{\sigma_k}), \quad (2.6)$$

where on the right hand side, $\mathbf{c}_1(\theta_j)$ denotes Clifford multiplication by the vector field which corresponds to θ_j via the Riemannian structure on X . Each $\mathbf{c}(\theta_1 \wedge \cdots \wedge \theta_k)$ acts as multiplication operator on $L^2(X, \Sigma)$. The properties (2.2) are well-known to hold in this case. Moreover, (A2) is satisfied because \mathbf{D}^2 is a non-negative elliptic differential operator which thus has a smooth heat kernel; (A1) follows from elliptic estimates, as $[\mathbf{D}, \mathbf{c}(\theta)]$ is a first order differential operator for each θ . Note that

$$\mathbf{c} : \Omega(X) \longrightarrow \mathcal{L}(L^2(X, \Sigma))$$

is *not* multiplicative (in particular, not a representation), if $\dim(X) \geq 2$.

This example has obvious generalizations to general unitary Clifford modules in the sense of [7, Def. 3.32].

Example 2.4 (Spin manifolds over X). Generalizing the previous example, if X is any manifold and E is a vector bundle with connection over X , then compact spin^c manifolds M over X give rise to a ϑ -summable Fredholm module $\mathcal{M} = (\mathcal{H}, \mathbf{c}, Q)$ over the locally convex algebra $\Omega(X)$. More precisely, suppose we are given an even-dimensional smooth spin^c manifold M with spinor bundle Σ_M , together with a smooth map $\varphi : M \rightarrow X$. Then we can set $\mathcal{H} = L^2(M, \Sigma_M \otimes \varphi^*E)$, $Q = \mathbf{D}_\varphi$ (the Dirac operator on M , twisted by φ^*E) and $\mathbf{c}(\theta) = \mathbf{c}_M(\varphi^*\theta)$, where \mathbf{c}_M is the quantization map on M as discussed in Example 2.3.

We now describe an important construction in this paper, which produces a new Fredholm module $\mathcal{M}_{\mathbb{T}}$ from a given Fredholm module \mathcal{M} ; in fact, this will only be a *weak* Fredholm module, which is our main reason to considering this weaker notion in the first place. The motivation for this construction is its connection to equivariant homology on loop spaces in the special case of Example 2.3.

We will see in §7 that the Bismut-Chern character $\text{Ch}(p)$ of an idempotent p in Ω^0 is not a chain over Ω but rather a chain over the acyclic extension $\Omega_{\mathbb{T}}$; hence we are going to need the Chern character of $\mathcal{M}_{\mathbb{T}}$ in order to formulate our index theorem.

Example 2.5 (Acyclic extension of a Fredholm module). Given a locally convex dg algebra Ω , we can form the dg algebra

$$\Omega_{\mathbb{T}} := \Omega[\sigma],$$

where σ is a formal variable of degree -1 with $\sigma^2 = 0$. Elements $\theta \in \Omega_{\mathbb{T}}$ can be written uniquely in the form $\theta = \theta' + \sigma\theta''$, where $\theta', \theta'' \in \Omega$, and the differential is

$$d_{\mathbb{T}} = d - \iota \quad \text{with} \quad d\theta = d\theta' - \sigma d\theta'' \quad \text{and} \quad \iota\theta = \theta''. \quad (2.7)$$

Then $\Omega_{\mathbb{T}}$ becomes a locally convex dg algebra in view of the vector space isomorphism $\Omega_{\mathbb{T}} \cong \Omega \oplus \Omega[1]$, called the *acyclic extension* of Ω . $\Omega_{\mathbb{T}}$ is indeed acyclic, as all closed elements have the form $\theta' + \sigma d\theta' = -d_{\mathbb{T}}(\sigma\theta')$, hence are exact.

Now given a ϑ -summable (weak) Fredholm module $\mathcal{M} = (\mathcal{H}, \mathbf{c}, Q)$, the *acyclic extension* of \mathcal{M} is the ϑ -summable weak Fredholm module $\mathcal{M}_{\mathbb{T}} = (\mathcal{H}, \mathbf{c}_{\mathbb{T}}, Q)$ over $\Omega_{\mathbb{T}}$, where $\mathbf{c}_{\mathbb{T}}$ is given by

$$\mathbf{c}_{\mathbb{T}}(\theta' + \sigma\theta'') := \mathbf{c}(\theta').$$

We will often write \mathbf{c} again instead of $\mathbf{c}_{\mathbb{T}}$. Notice that usually, this is indeed only a weak Fredholm module, since $(\Omega_{\mathbb{T}})^0$ is strictly larger than Ω^0 unless $\Omega^1 = \{0\}$.

Remark 2.6. In the special case that $\Omega = \Omega(X)$, sending σ to dt gives an isomorphism of algebras

$$\Omega_{\mathbb{T}} \cong \Omega(X \times S^1)^{S^1}, \quad (2.8)$$

the algebra of differential forms on $X \times S^1$ that are invariant under the S^1 -action on the second factor. Under this isomorphism, the differential ι on $\Omega_{\mathbb{T}}$ corresponds to the insertion of the vector field ∂_t on S^1 . Note that the isomorphism (2.8) does not preserve degrees, as we declared σ to have degree -1 in order to make the differential $d_{\mathbb{T}}$ homogeneous. Alternatively, this can be fixed by introducing a formal variable of degree two, but we shall avoid that in this paper.

3 Algebraic Preliminaries

In this section, we discuss the algebraic preliminaries needed for the construction of our Chern character. In particular, we introduce the chain and cochain complexes that we will be working on: The bar complex and the cyclic complex, as well as its entire and Chen normalized versions. Throughout this section, let Ω be a dg algebra.

3.1 Bar Chains and Cochains

The *bar construction* on Ω is the graded vector space

$$\mathbf{B}(\Omega) = \bigoplus_{N=0}^{\infty} \Omega[1]^{\otimes N}. \quad (3.1)$$

Here $\Omega[1]$ denotes the \mathbb{Z} -graded space with $\Omega[1]^k = \Omega^{k+1}$. The grading on $\mathbf{B}(\Omega)$ is then explicitly given by

$$\mathbf{B}_n(\Omega) = \bigoplus_{N=0}^{\infty} \bigoplus_{\ell_1 + \dots + \ell_N = n+N} \Omega^{\ell_1} \otimes \dots \otimes \Omega^{\ell_N}.$$

Elements of $\mathbf{B}(\Omega)$ will be called *bar chains on Ω* and denoted by $(\theta_1, \dots, \theta_N)$, omitting the tensor product sign in notation. The space $\mathbf{B}(\Omega)$ carries the two differentials d and b' ,

given by⁵

$$d(\theta_1, \dots, \theta_N) = \sum_{k=1}^N (-1)^{n_{k-1}} (\theta_1, \dots, \theta_{k-1}, d\theta_k, \dots, \theta_N)$$

$$b'(\theta_1, \dots, \theta_N) = - \sum_{k=1}^{N-1} (-1)^{n_k} (\theta_1, \dots, \theta_{k-1}, \theta_k \theta_{k+1}, \theta_{k+2}, \dots, \theta_N)$$

where $n_k = |\theta_1| + \dots + |\theta_k| - k$. The above differentials satisfy $db' + b'd = 0$, hence turn $\mathbf{B}(\Omega)$ into a bi-complex with total differential $d + b'$.

Given a \mathbb{Z}_2 -graded algebra L of coefficients, we define an L -valued bar cochain on Ω to be an element of the space $\text{Hom}(\mathbf{B}(\Omega), L)$ of linear maps from $\mathbf{B}(\Omega)$ to L . Such an L -valued bar cochain can be identified with a sequence $\ell^{(0)}, \ell^{(1)}, \ell^{(2)}, \dots$, consisting of its arity N components; each component is a multi-linear map

$$\ell^{(N)} : \underbrace{\Omega \times \dots \times \Omega}_N \longrightarrow L.$$

In particular, the component $\ell^{(0)}$ is a linear map from \mathbb{C} to L , which can be identified with an element of L .

Following Quillen [47] we introduce a sign in the definition of the *codifferential*

$$\delta := -(d + b') : \text{Hom}(\mathbf{B}(\Omega), L) \longrightarrow \text{Hom}(\mathbf{B}(\Omega), L), \quad (3.2)$$

which is defined as in (1.7). It will be important for our constructions that the space of L -valued bar cochains is a \mathbb{Z}_2 -graded algebra with respect to the product given by

$$\ell_1 \ell_2(\theta_1, \dots, \theta_N) = \sum_{k=1}^N (-1)^{|\ell_2|(|\theta_1| + \dots + |\theta_k|)} \ell_1(\theta_1, \dots, \theta_k) \ell_2(\theta_{k+1}, \dots, \theta_N), \quad (3.3)$$

in a way that δ satisfies the \mathbb{Z}_2 -graded Leibniz rule

$$\delta(\ell_1 \ell_2) = (\delta \ell_1) \ell_2 + (-1)^{|\ell_1|} \ell_1 (\delta \ell_2).$$

Thus δ turns $\text{Hom}(\mathbf{B}(\Omega), L)$ into a differential \mathbb{Z}_2 -graded algebra.

3.2 Cyclic Chains and Cochains

Throughout, we denote by $\underline{\Omega}$ the quotient space $\underline{\Omega} := \Omega / \mathbb{C}\mathbf{1}$. The (*reduced*) cyclic complex of Connes associated to Ω is the graded vector space

$$\mathbf{C}(\Omega) = \bigoplus_{N=0}^{\infty} \Omega \otimes \underline{\Omega}[1]^{\otimes N}, \quad (3.4)$$

⁵The sign of b' coincides with that of Quillen's b' [47], who has considered the case of ungraded algebras.

where we remind the reader that $\underline{\Omega}[1]$ denotes the graded vector space with $\underline{\Omega}[1]^k = \underline{\Omega}^{k+1}$. Explicitly, the grading of $\mathbf{C}(\Omega)$ is therefore given by

$$\mathbf{C}_n(\Omega) = \bigoplus_{M=0}^{\infty} \bigoplus_{\ell_0+\dots+\ell_M=n+M} \Omega^{\ell_0} \otimes \underline{\Omega}^{\ell_1} \otimes \dots \otimes \underline{\Omega}^{\ell_M}.$$

Elements of $\mathbf{C}(\Omega)$ will be called *cyclic chains*, denoted by $(\theta_0, \dots, \theta_N)$. The complex $\mathbf{C}(\Omega)$ carries two degree +1 differentials, given by the formulas

$$\begin{aligned} \underline{d}(\theta_0, \dots, \theta_N) &= (d\theta_0, \theta_1, \dots, \theta_N) - \sum_{k=1}^N (-1)^{m_{k-1}} (\theta_0, \dots, \theta_{k-1}, d\theta_k, \theta_{k+1}, \dots, \theta_N) \\ \underline{b}(\theta_0, \dots, \theta_N) &= \sum_{k=0}^{N-1} (-1)^{m_k} (\theta_0, \dots, \theta_k \theta_{k+1}, \dots, \theta_N) - (-1)^{(|\theta_N|-1)m_{N-1}} (\theta_N \theta_0, \theta_1, \dots, \theta_{N-1}). \end{aligned}$$

where $m_k = |\theta_0| + \dots + |\theta_k| - k$. It is easy to check that these formulas make sense on the quotient space $\Omega \otimes \underline{\Omega}[1]^{\otimes N}$. There is another differential, of degree -1 , the *Connes operator* \underline{B} , which is given by the formula

$$\underline{B}(\theta_0, \dots, \theta_N) = \sum_{k=0}^N (-1)^{(m_k+1)(m_N-m_k)} (\mathbf{1}, \theta_{k+1}, \dots, \theta_N, \theta_0, \dots, \theta_k)$$

with m_k as before. We remark that the Connes differential is one reason to restrict discussion to the reduced case, since its formula becomes more complicated otherwise, c.f. e.g. [34, p. 13]. The three differentials \underline{d} , \underline{b} and \underline{B} pairwise anti-commute, hence we get a \mathbb{Z}_2 -graded complex

$$\begin{array}{ccc} \mathbf{C}_+(\Omega) & \xrightarrow{\underline{d}+\underline{b}+\underline{B}} & \mathbf{C}_-(\Omega), \\ & \xleftarrow{\underline{d}+\underline{b}+\underline{B}} & \end{array} \quad (3.5)$$

where the spaces $\mathbf{C}_{\pm}(\Omega)$ are given by reducing the \mathbb{Z} -grading of $\mathbf{C}(\Omega)$ modulo two. Note that this is not a \mathbb{Z} -graded complex, as the differential is inhomogeneous.

3.3 Analytic Cochains and Entire Chains

In this section, let Ω be a locally convex dg algebra. As for bar cochains, a complex-valued cochain ℓ on $\mathbf{C}(\Omega)$ can be identified with the sequence $\ell^{(0)}, \ell^{(1)}, \ell^{(2)}, \dots$ of its arity N components, each component being a multi-linear map

$$\ell^{(N)} : \Omega \times \underbrace{\underline{\Omega}[1] \times \dots \times \underline{\Omega}[1]}_N \longrightarrow \mathbb{C}.$$

Following the nomenclature of [44, §2.1.2], we introduce the following growth condition on cochains.

Definition 3.1 (Analytic cochains). A cochain $\ell \in \text{Hom}(\mathbf{C}(\Omega), \mathbb{C})$ is *analytic* if there exists a continuous seminorm ν on Ω and a constant $C > 0$, such that⁶

$$|\ell^{(N)}(\theta_0, \dots, \theta_N)| \leq \frac{C}{[N/2]!} \nu(\theta_0) \cdots \nu(\theta_N) \quad (3.6)$$

for all $\theta_0, \dots, \theta_N \in \Omega$. The space of analytic cochains will be denoted by $\mathbf{C}_\alpha(\Omega)$.

In particular, each of the linear maps $\ell^{(N)}$ is continuous with respect to the projective tensor product topology on $\Omega \otimes \underline{\Omega}^{\otimes N}$, see below. The differentials \underline{d} , \underline{b} and \underline{B} are easily checked to preserve the space $\mathbf{C}_\alpha(\Omega)$, leading to the *analytic cocomplex* of Ω .

The growth conditions imposed the components $\ell^{(N)}$ of analytic cochains ℓ allow to evaluate such a cochain on certain chains that are an infinite sum of their homogenous tensor components, in other words, element of the direct *product* of the $\Omega \otimes \Omega[1]^{\otimes N}$. To define these *entire chains*, we recall that for each continuous seminorm ν on Ω the induced N -th *projective tensor norm* $\pi_{\nu, N}$ on $\Omega \otimes \Omega[1]^{\otimes N}$ is defined by the formula

$$\pi_{\nu, N}(c_N) = \inf \left\{ \sum_{\gamma} \nu(\theta_0^{(\gamma)}) \cdots \nu(\theta_N^{(\gamma)}) : c_N = \sum_{\gamma} \theta_0^{(\gamma)} \otimes \cdots \otimes \theta_N^{(\gamma)} \right\}, \quad (3.7)$$

where the infimum is taken over all representations of $c_N \in \mathbf{C}(\Omega)$ as a finite sum of elementary tensors.

Definition 3.2 (Entire chains). For each continuous seminorm ν on Ω , we define a seminorm ϵ_ν on $\mathbf{C}(\Omega)$ by

$$\epsilon_\nu(c) := \sum_{N=0}^{\infty} \frac{\pi_{\nu, N}(c_N)}{[N/2]!}, \quad c = \sum_{N=0}^{\infty} c_N \in \mathbf{C}(\Omega), \quad \text{with } c \in \Omega \otimes \Omega[1]^{\otimes N}, \quad (3.8)$$

which is finite as the definition of $\mathbf{C}(\Omega)$ involves the direct sum. The completion of $\mathbf{C}(\Omega)$ with respect to the seminorms ϵ_ν is denoted by $\mathbf{C}^\epsilon(\Omega)$. Its elements are called *entire chains*.

It is easy to see that analytic cochains $\ell \in \mathbf{C}_\alpha(\Omega)$ are bounded with respect to the seminorms ϵ_ν , hence they extend by continuity to functionals on $\mathbf{C}^\epsilon(\Omega)$. Moreover, as the differentials \underline{d} , \underline{b} , \underline{B} are easily checked to be continuous on $\mathbf{C}(\Omega)$ with respect to the seminorms ϵ_ν , they extend to odd parity differentials on $\mathbf{C}^\epsilon(\Omega)$; we denote these extensions by the same symbols again, leading to the *entire complex* of Ω .

Remark 3.3. The terminology *analytic* for the growth condition (3.6) stems from the fact that this condition implies that the function

$$f(\theta) = \sum_{N=0}^{\infty} [N/2]! \ell^{(N)}(\underbrace{\theta, \dots, \theta}_{N+1})$$

⁶Here $[N/2] = N/2$ if N is even, and $[N/2] = (N-1)/2$ if N is odd.

defines an analytic function on some small enough ν -ball around zero in Ω . Similarly, for any entire chain c and any continuous seminorm ν on Ω , the function

$$f(z) := \sum_{N=0}^{\infty} \pi_{\nu,N}(c_N) \frac{z^N}{[N/2]!}$$

is an entire function of z . The entire growth condition was first introduced by Connes [21] for *cochains* on the space $\mathbf{C}(\mathcal{A})$ for a Banach algebra \mathcal{A} ; it then turns out [44, §2.1.2] that the space of entire cochains is the dual to the space of chains satisfying the analytic growth condition.

It turns out that in our context, it is suitable to consider the entire growth condition on chains rather than on cochains, and the analytic growth condition on cochains, rather than on chains, due to the fact that the cochain Chern character $\text{Ch}(\mathcal{M})$ satisfies the analytic growth condition (c.f. Thm. 4.6), while the Chern character defined in Def. 7.2 satisfies the entire growth condition, but not the analytic one.

For the algebra of differential forms on a manifold, these growth conditions on chains were previously considered e.g. in [41] and [40]; see also [35].

3.4 Chen Normalization

Again, we assume that Ω is a locally convex dg algebra with acyclic extension $\Omega_{\mathbb{T}}$, discussed in Example 2.5. In this section, we construct the normalized complexes that our constructions will ultimately live on. To this end, we consider a certain subcomplex $\mathbf{D}_{\mathbb{T}}(\Omega)$ of $\mathbf{C}(\Omega_{\mathbb{T}})$.

To define the subcomplex $\mathbf{D}^{\mathbb{T}}(\Omega)$ of $\mathbf{C}(\Omega_{\mathbb{T}})$ consider for all $f \in \Omega^0$ and $i \in \mathbb{N}_0$ the continuous linear map

$$S_i^{(f)} : \mathbf{C}(\Omega_{\mathbb{T}}) \longrightarrow \mathbf{C}(\Omega_{\mathbb{T}})$$

defined by

$$S_i^{(f)}(\theta_0, \dots, \theta_N) = \begin{cases} (\theta_0, \dots, \theta_i, f, \theta_{i+1}, \dots, \theta_N) & 0 \leq i \leq N \\ 0 & \text{otherwise} \end{cases}. \quad (3.9)$$

We emphasize here that we only consider $f \in \Omega^0 \subset \Omega_{\mathbb{T}}^0$, which is generally a proper subset. Consider further the maps

$$S : \mathbf{C}(\Omega_{\mathbb{T}}) \longrightarrow \mathbf{C}(\Omega_{\mathbb{T}}), \quad S(\theta_0, \dots, \theta_N) = \sum_{k=0}^N (\theta_0, \dots, \theta_k, \sigma, \theta_{k+1}, \dots, \theta_N), \quad (3.10)$$

$$R : \mathbf{C}(\Omega_{\mathbb{T}}) \longrightarrow \mathbf{C}(\Omega_{\mathbb{T}}), \quad R(\theta_0, \dots, \theta_N) = (\sigma\theta_0, \dots, \theta_N). \quad (3.11)$$

We then define the subcomplex $\mathbf{D}^{\mathbb{T}}(\Omega) \subset \mathbf{C}(\Omega_{\mathbb{T}})$ by

$$\mathbf{D}^{\mathbb{T}}(\Omega) := \left\{ \begin{array}{l} \text{The span of the images of the operators} \\ S, R \text{ and } S_i^{(f)}, [\underline{d} + \underline{b}, S_i^{(f)}] \text{ for } f \in \Omega^0 \end{array} \right\}. \quad (3.12)$$

It is straightforward to check that the differentials \underline{d} , \underline{b} and \underline{B} preserve this complex.

Definition 3.4 (The Chen normalized complex). The *Chen normalized entire complex* is the quotient complex

$$\mathbf{N}^{\mathbb{T},\epsilon}(\Omega) := \mathbf{C}^\epsilon(\Omega_{\mathbb{T}}) / \overline{\mathbf{D}^{\mathbb{T}}(\Omega)}, \quad (3.13)$$

where we take the closure inside $\mathbf{C}^\epsilon(\Omega_{\mathbb{T}})$. The space of *extended Chen normalized analytic cochains* is the space $\mathbf{N}_{\mathbb{T},\alpha}(\Omega)$ of analytic cochains vanishing on $\mathbf{D}^{\mathbb{T}}(\Omega)$. We obtain the short complexes

$$\begin{array}{ccc} \mathbf{N}_+^{\mathbb{T},\epsilon}(\Omega) & \xrightarrow{\underline{d}+\underline{b}+\underline{B}} & \mathbf{N}_-^{\mathbb{T},\epsilon}(\Omega), \\ & \xleftarrow{\underline{d}+\underline{b}+\underline{B}} & \\ \mathbf{N}_{\mathbb{T},\alpha}^+(\Omega) & \xrightarrow{\underline{d}+\underline{b}+\underline{B}} & \mathbf{N}_{\mathbb{T},\alpha}^-(\Omega). \\ & \xleftarrow{\underline{d}+\underline{b}+\underline{B}} & \end{array} \quad \text{and}$$

Remark 3.5. The subcomplex $\mathbf{D}(\Omega)$ of $\mathbf{C}(\Omega)$, generated by the images of the operators $S_i^{(f)}$ and $[\underline{d} + \underline{b}, S_i^{(f)}]$, was first considered by Chen [20], in the case that $\Omega = \Omega(X)$ for a manifold X . He also showed that $\mathbf{D}(\Omega(X))$ is acyclic with respect to the differential $\underline{d} + \underline{b}$ in the case that X is connected, meaning that the projection from $\mathbf{C}(\Omega(X))$ to the quotient complex

$$\mathbf{N}(\Omega(X)) := \mathbf{C}(\Omega(X)) / \mathbf{D}(\Omega(X))$$

is a homotopy equivalence in this case (with respect to the differential $\underline{d} + \underline{b}$). This was suitably generalized by Getzler and Jones to general dg algebras [33, Section 5].

This quotient (3.13) was first considered by Getzler, Jones and Petrack in the special case $\Omega = \Omega(X)$ for a manifold X ; in fact, the image of $S - \text{id}$ can be written as an ideal with respect to the shuffle product, as defined in §4 of [34]; it is generated by the chain $(1, \sigma) - (1) \in \mathbf{C}(\Omega)$. Cast in these terms, such a quotient is considered on p. 28 of [34]. Clearly, there is a map of chain complexes

$$\mathbf{N}^\epsilon(\Omega) := \mathbf{C}^\epsilon(\Omega) / \overline{\mathbf{D}(\Omega)} \longrightarrow \mathbf{N}^{\mathbb{T},\epsilon}(\Omega),$$

natural in Ω . The results of Getzler, Jones and Petrack seem to indicate that one might suspect this map to be a quasi-isomorphism, at least in the special case $\Omega = \Omega(X)$.

4 The Quantization Map

In this section, we start with the construction of the Chern character for a ϑ -summable weak Fredholm module $\mathcal{M} := (\mathcal{H}, \mathbf{c}, Q)$ over a locally convex dg algebra, which we assume fixed throughout this section.

As observed by Quillen [47], an odd bar cochain ω with values in a \mathbb{Z}_2 -graded algebra L of coefficients can be seen as a *connection form* on the space of L -valued bar cochains, as we can consider the *connection*⁷ $\nabla := \delta + \omega$. Its *curvature* is given by

$$F := \nabla^2 = \delta\omega + \omega^2. \quad (4.1)$$

⁷Since ω does not necessarily have arity one, this should rather be called a *superconnection* in the sense of Quillen.

Note that since $\delta^2 = 0$, F is an L -valued bar cochain itself, as opposed to merely an operator acting on cochains. It satisfies the *Bianchi identity*

$$0 = [\nabla, F] = \delta F + [\omega, F], \quad (4.2)$$

since $[\nabla, F] = \nabla^3 - \nabla^3 = 0$. These two formulas lie at the heart of our investigations.

To apply this, we observe that the Fredholm module \mathcal{M} naturally provides such a connection form $\omega_{\mathcal{M}}$ with values in the algebra of operators on \mathcal{H} . It is defined by

$$\omega_{\mathcal{M}}^{(0)} = -Q, \quad \omega_{\mathcal{M}}^{(1)}(\theta) = \mathbf{c}(\theta), \quad \omega_{\mathcal{M}}^{(N)}(\theta_1, \dots, \theta_N) = 0, \quad N \geq 2, \quad (4.3)$$

and it is odd due to the grading shift in the definition of $\mathbf{B}(\Omega)$. Let

$$F_{\mathcal{M}} = \delta\omega_{\mathcal{M}} + \omega_{\mathcal{M}}^2$$

be the curvature of the connection ω . Explicitly, the components of $F_{\mathcal{M}}$ can be easily worked out to be given by

$$\begin{aligned} F_{\mathcal{M}}^{(0)} &= Q^2, \\ F_{\mathcal{M}}^{(1)}(\theta) &= \mathbf{c}(d\theta) - [Q, \mathbf{c}(\theta)] \\ F_{\mathcal{M}}^{(2)}(\theta_1, \theta_2) &= (-1)^{|\theta_1|} (\mathbf{c}(\theta_1\theta_2) - \mathbf{c}(\theta_1)\mathbf{c}(\theta_2)). \end{aligned} \quad (4.4)$$

with all higher components zero. Notice that the first component essentially is the curvature of the derivation on $\mathcal{L}(\mathcal{H})$ given by taking the commutator with Q , the second component measures the failure of \mathbf{c} and Q to satisfy the relation $\mathbf{c}(d\theta) = [Q, \mathbf{c}(\theta)]$ for $\theta \in \Omega$, while the third component measures the failure of \mathbf{c} to be multiplicative. What causes complications is that $\omega_{\mathcal{M}}$ does not take values in bounded operators (hence does not actually take values in an algebra). For this reason, Quillen's constructions are more of a guiding principle, and we have to put in some effort to make them work.

In order to define the Chern character of our Fredholm module, we now aim to define a bar cochain $\Phi_T^{\mathcal{M}}$ with values in $\mathcal{L}(\mathcal{H})$, the *quantization map*, by exponentiating the curvature $F_{\mathcal{M}}$ in the algebra $\text{Hom}(\mathbf{B}(\Omega), \mathcal{L}(\mathcal{H}))$, i.e. we would like to set

$$\Phi_T^{\mathcal{M}} \stackrel{\text{formally}}{=} e^{-TF_{\mathcal{M}}}, \quad (4.5)$$

where $T > 0$ is any real parameter. If Q was a bounded operator, this would be well-defined using the exponential series; however, the fact that Q is unbounded leads to analytical difficulties. To overcome these difficulties, the plan is to split

$$F_{\mathcal{M}} = Q^2 + F_{\mathcal{M}}^{\geq 1}, \quad (4.6)$$

where $F_{\mathcal{M}}^{\geq 1}$ is the part containing the cochains of arity at least one (which takes values in bounded operators on \mathcal{H}). Using this, we have (formally)

$$e^{-TF_{\mathcal{M}}} = e^{-T(Q^2 + F_{\mathcal{M}}^{\geq 1})}$$

which can be rewritten into a perturbation series with respect to the heat operator e^{-TQ^2} . To facilitate this, we first introduce the following notation.

Notation 4.1 (Bracket). Given suitable⁸ operators A_1, \dots, A_N on \mathcal{H} , we set

$$\{A_1, \dots, A_N\}_Q := \int_{\Delta_N} e^{-\tau_1 Q^2} A_1 e^{-(\tau_2 - \tau_1) Q^2} A_2 \dots e^{-(\tau_N - \tau_{N-1}) Q^2} A_N e^{-(1 - \tau_N) Q^2} d\tau \quad (4.7)$$

where $\Delta_N = \{\tau \in \mathbb{R}^n \mid 0 \leq \tau_1 \leq \dots \leq \tau_N \leq 1\}$ is the standard simplex. If there is no danger of confusion, we drop the subscript Q in notation.

In order to define $\Phi_T^{\mathfrak{M}}$, we will have to allow unbounded operators A_j to be put into the bracket above. The following fundamental estimate will facilitate this.

Lemma 4.2. *For $0 \leq a_0, \dots, a_N < 1$ let A_0, \dots, A_N be densely defined operators on \mathcal{H} such that $A_j \Delta^{-a_j}$ is densely defined and extends to a bounded operator on \mathcal{H} , where $\Delta = Q^2 + 1$. Then for each $T > 0$, $A_0 \{A_1, \dots, A_N\}_{TQ}$ is a well-defined trace-class operator, and we have the estimate*

$$\|A_0 \{A_1, \dots, A_N\}_{TQ}\|_1 \leq \frac{e^{T/2} \operatorname{Tr}(e^{-TQ^2/2})}{\Gamma(N + 1 - (a_0 + \dots + a_N))} \prod_{j=0}^N \|A_j \Delta^{-a_j}\| \left(\frac{2a_j \Gamma(1 - a_j)}{eT} \right)^{a_j},$$

for its nuclear norm, where Γ denotes the gamma function.

Proof. By the assumptions on A_j , for each $\varepsilon > 0$, the operator

$$A_j e^{-\varepsilon Q^2} = (A_j \Delta^{-a_j}) (\Delta^{a_j} e^{-\varepsilon/2 Q^2}) e^{-\varepsilon/2 Q^2}$$

is trace-class, hence for each fixed $\tau \in \Delta_N$ such that $\tau_j - \tau_{j-1} > 0$, the integrand in (4.7) is a trace-class operator, by the assumptions on the A_j . In order to show that the integral over Δ_N gives a well-defined trace-class operator, it then suffices to show that the nuclear norm as a function of τ is integrable.

To this end, set $s_j = \tau_{j+1} - \tau_j$, $j = 0, \dots, N$, where $\tau_{N+1} = 1$, $\tau_0 = 0$. Since $\sum_{j=0}^N s_j = 1$, we have using Hölder's inequality for Schatten norms

$$\begin{aligned} \left\| \prod_{j=0}^N A_j e^{-T(\tau_j - \tau_{j-1}) Q^2} \right\|_1 &= \left\| \prod_{j=0}^N A_j e^{-T s_j Q^2} \right\|_1 \leq \prod_{j=0}^N \|A_j \Delta^{-a_j}\| \|\Delta^{a_j} e^{-T s_j Q^2}\|_{1/s_j} \\ &\leq \prod_{j=0}^N \|A_j \Delta^{-a_j}\| \prod_{j=0}^N \|\Delta^{a_j} e^{-T s_j Q^2/2}\| \prod_{j=0}^N \|e^{-T s_j Q^2/2}\|_{1/s_j}. \end{aligned}$$

Now for the last product, we get

$$\prod_{j=0}^N \|e^{-T s_j Q^2/2}\|_{1/s_j} = \prod_{j=0}^N \left(\sum_{k=1}^{\infty} (e^{-T s_j \lambda_j/2})^{1/s_j} \right)^{s_j} = \prod_{j=0}^N \operatorname{Tr}(e^{-TQ^2/2})^{s_j} = \operatorname{Tr}(e^{-TQ^2/2}),$$

⁸In the sense that they satisfy the assumptions of Lemma 4.2 below.

where λ_j , for $j = 1, 2, \dots$, are the eigenvalues of Q^2 . To estimate the second product, observe that

$$(x^2 + 1)^a e^{-\tau x^2/2} \leq \left(\frac{2a}{s}\right)^a e^{\tau/2-a}.$$

for all $x \in \mathbb{R}$. Hence

$$\prod_{j=0}^N \|\Delta^{a_j} e^{-T s_j Q^2/2}\| \leq e^{T/2} \prod_{j=0}^N \left(\frac{2a_j}{eT}\right)^{a_j} (\tau_{j+1} - \tau_j)^{-a_j}$$

and the lemma now follows from the formula

$$\int_{\Delta_N} \tau_1^{-a_0} (\tau_2 - \tau_1)^{-a_1} \dots (1 - \tau_N)^{-a_N} d\tau = \frac{\Gamma(1 - a_0) \dots \Gamma(1 - a_N)}{\Gamma(N + 1 - (a_0 + \dots + a_N))};$$

in the case $N = 1$, this latter formula is the usual integral formula for the beta function, while the case for general N can be easily shown by induction. \square

As can be seen from 4.4, the operators $F_{\mathcal{M}}(\theta)$ are unbounded operators in general. However, by the assumption (A1) of the Fredholm module, they satisfy the assumptions of Lemma 4.2. Therefore, using the algebra structure on the space of $\mathcal{L}(\mathcal{H})$ -valued bar cochains, the expression $\{F_{\mathcal{M}}^{\geq 1}, \dots, F_{\mathcal{M}}^{\geq 1}\}_{TQ}$, used in the definition below, defines a well-defined cochain.

Definition 4.3 (Quantization Map). For $T > 0$ fixed, the *quantization map* $\Phi_T^{\mathcal{M}}$ is the $\mathcal{L}(\mathcal{H})$ -valued bar cochain defined by the formula

$$\Phi_T^{\mathcal{M}} := \sum_{N=0}^{\infty} (-T)^N \underbrace{\{F_{\mathcal{M}}^{\geq 1}, \dots, F_{\mathcal{M}}^{\geq 1}\}}_N{}_{TQ}. \quad (4.8)$$

Remark 4.4. $\Phi_T^{\mathcal{M}}$ is an even cochain, in the sense that it maps even elements of $\mathbf{B}(\Omega)$ to even operators on \mathcal{H} and odd elements to odd operators. This follows from the fact that F has this property (which in turn holds because ω is odd, or can be seen from an inspection of (4.4) above), and the fact that the bracket is also even in the obvious sense (as it depends only on the square of Q).

To state explicitly how Φ_T evaluates on chains, we need the following notation.

Notation 4.5 (Partitions). Denote by $\mathcal{P}_{M,N}$ the set of ordered partitions of the set $\{1, \dots, N\}$ of length M , i.e. collections of nonempty subsets $I = (I_1, \dots, I_M)$ such that $I_1 \cup \dots \cup I_M = \{1, \dots, N\}$ and such that each element of I_a is smaller than any element of I_b whenever $a < b$.

Now an inspection of the definition of $\Phi_T^{\mathcal{M}}$ gives the formula

$$\Phi_T^{\mathcal{M}}(\theta_1, \dots, \theta_N) = \sum_{M=1}^N (-T)^M \sum_{I \in \mathcal{P}_{M,N}} \{F_{\mathcal{M}}(\theta_{I_1}), \dots, F_{\mathcal{M}}(\theta_{I_M})\}_{TQ}, \quad (4.9)$$

where we used the obvious notation $\theta_{I_a} := (\theta_{i+1}, \dots, \theta_{i+m})$ if $I_a = \{j \mid i < j \leq i+m\}$ for some i, m . This formula follows from the product formula (3.3) for cochains. Notice that a summand corresponding to $I = (I_1, \dots, I_M) \in \mathcal{P}_{M,N}$ is zero as soon as I contains an I_a with $|I_a| \geq 3$.

The following are now the main results of this section.

Theorem 4.6 (Fundamental estimate). *For $T > 0$ and all $\theta_1, \dots, \theta_N \in \Omega$, the operator $\Phi_T^{\mathcal{M}}(\theta_1, \dots, \theta_N)$ is well-defined and trace-class. Moreover, there exists a continuous seminorm ν on Ω , independent of T , such that we have the estimate*

$$\|\Phi_T^{\mathcal{M}}(\theta_1, \dots, \theta_N)\|_1 \leq e^{T/2} \text{Tr}(e^{-TQ^2/2}) \frac{T^N}{[n/2]!} \nu(\theta_1) \cdots \nu(\theta_N) \quad (4.10)$$

for its trace class norm. The same statement is true for $Q\Phi_T^{\mathcal{M}}$ and $\Phi_T^{\mathcal{M}}Q$ instead of $\Phi_T^{\mathcal{M}}$.

Theorem 4.7 (Bianchi identity). *The map $\Phi_T^{\mathcal{M}}$ satisfies*

$$\delta\Phi_T^{\mathcal{M}} + [\omega, \Phi_T^{\mathcal{M}}] = 0.$$

Thm. 4.7 is obvious at a formal level, using the formal equality (4.5) together with the Bianchi identity (4.2) for $F_{\mathcal{M}}$, but again, the fact that $F_{\mathcal{M}}$ does not take values in $\mathcal{L}(\mathcal{H})$ does not allow this easy proof.

Remark 4.8. Explicitly, on elements, this identity can be worked out to state

$$\begin{aligned} \Phi_T^{\mathcal{M}}((d+b')(\theta_1, \dots, \theta_N)) &= \mathbf{c}(\theta_1)\Phi_T^{\mathcal{M}}(\theta_2, \dots, \theta_N) - (-1)^{n_{N-1}}\Phi_T^{\mathcal{M}}(\theta_1, \dots, \theta_{N-1})\mathbf{c}(\theta_N) \\ &\quad - [Q, \Phi_T^{\mathcal{M}}(\theta_1, \dots, \theta_N)], \end{aligned} \quad (4.11)$$

where $n_k = |\theta_1| + \dots + |\theta_k| - k$. Here both sides are well-defined by Thm. 4.6.

Proof (of Thm. 4.6). We drop the dependence on \mathcal{M} in the notation. First consider the cochain Φ_T . The operator $\Phi_T(\theta_1, \dots, \theta_N)$ is a sum of operators $A_0\{A_1, \dots, A_N\}_{TQ}$, with $A_0 = 0$ and each A_j , $j \geq 1$ equal to one of the terms in (4.4). Now by the properties (2.1) of the locally convex topology and the assumptions (A2) on the Fredholm module, there exists a continuous seminorm ν on Ω such that

$$\|F(\theta)\Delta^{-1/2}\| \leq \nu(\theta) \quad \text{and} \quad \|F(\theta_1, \theta_2)\| \leq \nu(\theta_1)\nu(\theta_2). \quad (4.12)$$

Hence Lemma 4.2 implies is a well-defined trace-class operator for each N . The same is true for $Q\Phi_T(\theta_1, \dots, \theta_N)$, only that in this case, $A_0 = Q$. For $\Phi_T(\theta_1, \dots, \theta_N)Q$, we can argue by passing to the adjoint.

It therefore remains to show the estimate (4.10). Here we consider the cochain Φ_T , the proof for $Q\Phi_T$ and $\Phi_T Q$ is similar. Now notice that since F has only components of arity less than two, for any partition $I \in \mathcal{P}_{M,N}$, we have

$$\{F(\theta_{I_1}), \dots, F(\theta_{I_M})\}_{TQ} = 0$$

as soon as I contains a subset I_j with $|I_j| \geq 2$. For all other partitions, we have $|I_j| = 1$ for $2M - N$ indices j and $|I_j| = 2$ for $N - M$ indices j . Hence by (4.12), we can apply Lemma 4.2 with $a_j = 1/2$ for $2M - N$ and $a_j = 0$ for $N - M$ indices j . This gives

$$\|\{F(\theta_{I_1}), \dots, F(\theta_{I_M})\}_T\|_1 \leq \frac{e^{T/2} \operatorname{Tr}(e^{-TQ^2/2})}{\Gamma(\frac{3}{2}N + 1 - M)} \left(\frac{\sqrt{\pi}}{eT}\right)^{(2M-N)/2} \nu(\theta_1) \cdots \nu(\theta_N),$$

as $\Gamma(1/2) = \sqrt{\pi}$. With a view on (4.9), we therefore get

$$\begin{aligned} \|\Phi_T(\theta_1, \dots, \theta_N)\|_1 &\leq \sum_{M=\lceil N/2 \rceil}^N T^M \sum_{I \in \mathcal{P}_{M,N}} \frac{e^{T/2} \operatorname{Tr}(e^{-TQ^2/2})}{\Gamma(\frac{3}{2}N + 1 - M)} \left(\frac{\sqrt{\pi}}{eT}\right)^{(2M-N)/2} \nu(\theta_1) \cdots \nu(\theta_N) \\ &\leq T^{N/2} \frac{e^{T/2} \operatorname{Tr}(e^{-TQ^2/2})}{\Gamma(N/2)} \sum_{M=\lceil N/2 \rceil}^N \sum_{I \in \mathcal{P}_{M,N}} \left(\frac{\sqrt{\pi}}{e}\right)^{(2M-N)/2} \nu(\theta_1) \cdots \nu(\theta_N) \end{aligned}$$

The sum over M growth at most like C^N with N , where $C > 0$ is some constant, hence it can be absorbed into the seminorm (by replacing it by a larger one). \square

Proof (of Thm. 4.7). Again we drop the dependence on \mathcal{M} in the notation. For the proof, we introduce for $\varepsilon \geq 0$ the Bracket $\{A_1, \dots, A_N\}_Q^\varepsilon$, which is defined by the same formula (4.7), integrating over

$$\Delta_N^\varepsilon = \{\tau \in \Delta_N \mid \tau_j - \tau_{j-1} \geq \varepsilon \text{ for all } j\}$$

but instead of Δ_N , the N -simplex with the ε -neighborhood of the diagonals removed. Due to the fact that for any m , the norm of the operator $Q^k e^{-tQ^2}$ is bounded uniformly on $[\varepsilon, 1]$, the expression $\{A_1, \dots, A_N\}_Q^\varepsilon$ gives a well-defined operator for any collection of operators A_j of the form $A_j = Q^n A'_j Q^m$, with A'_j bounded. We denote by Φ_T^ε the cochain defined by the same formula (4.8) as Φ_T , but using this bracket instead of the previous one. Now for any $\varepsilon \geq 0$, we have

$$\delta\Phi_T^\varepsilon = \sum_{N=0}^{\infty} (-T)^N \sum_{k=1}^N \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k}, \delta F^{\geq 1}, \underbrace{F^{\geq 1}, \dots, F^{\geq 1}}_{k-1} \Big\}_{TQ}^\varepsilon. \quad (4.13)$$

By the Bianchi identity (4.2) for F , we have

$$\delta F^{\geq 1} = \delta F = -[\omega, F] = [Q, F^{\geq 1}] - [\mathbf{c}, F^{\geq 1}] - [\mathbf{c}, Q^2]. \quad (4.14)$$

At this point, this identity is just formal as both side are cochains with values in unbounded operators and we do not have enough control over their domains; however, it is easy to verify that

$$\begin{aligned}
\delta\Phi_T^\varepsilon &= \sum_{N=1}^{\infty} (-T)^N \sum_{k=1}^N \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1} \delta F^{\geq 1} \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k}^\varepsilon_{TQ} \\
&= \sum_{N=1}^{\infty} (-T)^N \left(\sum_{k=1}^N \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1} [Q, F^{\geq 1}] \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k}^\varepsilon_{TQ} \right. \\
&\quad - \sum_{k=1}^N \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1} [\mathbf{c}, F^{\geq 1}] \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k}^\varepsilon_{TQ} \\
&\quad \left. - \sum_{k=1}^N \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1} [\mathbf{c}, Q^2] \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k}^\varepsilon_{TQ} \right), \tag{4.15}
\end{aligned}$$

formally obtained by plugging the right hand of (4.14) side into (4.13), is in fact an identity of well-defined cochains provided that $\varepsilon > 0$, by the discussion of the ε -bracket above. Notice that since taking the commutator with Q is a derivation, we have

$$\underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1} [Q, F^{\geq 1}] \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k}^\varepsilon_{TQ} = \left[Q, \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_N^\varepsilon_{TQ} \right],$$

which in fact makes sense even for $\varepsilon = 0$, by Lemma 4.6 above. Also the second term on the right hand side of (4.15) makes sense for $\varepsilon = 0$; for the third term, we use the following lemma.

Lemma 4.9. *Let A_1, \dots, A_N be a collection of operators of the form $A_j = Q^m A'_j Q^n$ with A'_j bounded and assume that $A_k = A'_k$ for some k . Then for $\varepsilon > 0$, we have the identity*

$$\begin{aligned}
&T\{A_1, \dots, [Q^2, A_k], \dots, A_N\}^\varepsilon_{TQ} \\
&= \{A_1, \dots, A_k e^{-\varepsilon Q^2} A_{k+1}, \dots, A_N\}^\varepsilon_{TQ} - \{A_1, \dots, A_{k-1} e^{-\varepsilon Q^2} A_k, \dots, A_N\}^\varepsilon_{TQ}.
\end{aligned}$$

Here, if $k = 1$, the second term on the right-hand-side is $A_1\{A_2, \dots, A_N\}^\varepsilon_{TQ}$ instead, while if $k = N$, the first term is $\{A_1, \dots, A_{N-1}\}^\varepsilon_T A_N$.

Proof. This follows from integration by parts, after realizing that the term on the left hand side equals

$$\int_{\Delta_N^\varepsilon} \frac{\partial}{\partial \tau_k} \left[e^{-T(1-\tau_N)Q^2} \prod_{j=1}^N A_j e^{-T(\tau_j - \tau_{j-1})Q^2} \right] d\tau. \quad \square$$

Hence for any $\varepsilon > 0$, we have

$$\begin{aligned} \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1}, [\mathbf{c}, Q^2], \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k} \Big|_{TQ} &= \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1}, \mathbf{c}e^{-\varepsilon Q^2} F^{\geq 1}, \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k-1} \Big|_{TQ} \\ &\quad - \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-2}, F^{\geq 1} e^{-\varepsilon Q^2} \mathbf{c}, \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k} \Big|_{TQ}. \end{aligned}$$

Now notice that the right hand side of the above identity makes sense for $\varepsilon = 0$ as well. Hence after replacing the last term in (4.15) with this term, we may take the limit $\varepsilon \rightarrow 0$ in the identity (4.15). This gives

$$\begin{aligned} \delta\Phi_T &= [Q, \Phi_T] - \sum_{N=1}^{\infty} (-T)^N \sum_{k=1}^N \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1}, [\mathbf{c}, F^{\geq 1}], \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k} \Big|_{TQ} \\ &\quad - \sum_{N=1}^{\infty} (-T)^{N-1} \left(\mathbf{c} \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-1} \Big|_{TQ} - \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-1} \Big|_{TQ} \mathbf{c} \right. \\ &\quad \quad \quad + \sum_{k=2}^N \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-2}, F^{\geq 1} \mathbf{c}, \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k} \Big|_{TQ} \\ &\quad \quad \quad \left. - \sum_{k=1}^{N-1} \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1}, \mathbf{c} F^{\geq 1}, \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k-1} \Big|_{TQ} \right). \end{aligned}$$

Finally, after an index shift, we obtain that the third term on the right hand side equals

$$\begin{aligned} & - \sum_{N=1}^{\infty} (-T)^{N-1} \left(\mathbf{c} \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-1} \Big|_T - \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-1} \Big|_{TQ} \mathbf{c} \right. \\ & \quad \quad \quad \left. + \sum_{k=1}^{N-1} \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1}, F^{k \geq 1} \mathbf{c} - \mathbf{c} F^{\geq 1}, \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k-1} \Big|_{TQ} \right) \\ &= -[\mathbf{c}, \Phi_T] + \sum_{N=2}^{\infty} (-T)^{N-1} \sum_{k=1}^{N-1} \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{k-1}, [\mathbf{c}, F^{\geq 1}], \underbrace{\{F^{\geq 1}, \dots, F^{\geq 1}\}}_{N-k-1} \Big|_{TQ} \end{aligned}$$

Combining this with the identity before, we obtain

$$\delta\Phi_T = [Q, \Phi_T] - [\mathbf{c}, \Phi_T] = -[\omega, \Phi_T],$$

which was the claim. \square

5 The Chern Character

Throughout this section, let Ω be a locally convex dg algebra.

Definition 5.1 (Chern character). For any ϑ -summable weak Fredholm module \mathcal{M} , we define an even cochain $\text{Ch}(\mathcal{M}) \in \text{Hom}^+(\mathbb{C}(\Omega), \mathbb{C})$ by

$$\langle \text{Ch}(\mathcal{M}), (\theta_0, \dots, \theta_N) \rangle = \text{Str} \left(\mathbf{c}(\theta_0) \Phi_1^{\mathcal{M}}(\theta_1, \dots, \theta_N) \right)$$

for $\theta_0, \dots, \theta_N \in \Omega$. Here $\Phi_T^{\mathcal{M}}$ is the quantization map defined in §4 and Str denotes the supertrace for trace-class operators on the \mathbb{Z}_2 -graded Hilbert space \mathcal{H} . The cochain $\text{Ch}(\mathcal{M})$ is called the *Chern character* of \mathcal{M} .

Note that $\text{Ch}(\mathcal{M})$ is well-defined: First, that the operator $\mathbf{c}(\theta_0) \Phi_1^{\mathcal{M}}(\theta_1, \dots, \theta_N)$ is trace-class follows from the fundamental estimate from Thm. 4.6. Secondly, $\text{Ch}(\mathcal{M})$ is a well-defined element of $\text{Hom}(\mathbb{C}(\Omega), \mathbb{C})$. To see this, one has to observe that

$$\Phi_1^{\mathcal{M}}(\theta_1, \dots, \theta_j, \mathbf{1}, \theta_{j+1}, \dots, \theta_N) = 0$$

for all $\theta_1, \dots, \theta_N \in \Omega$ and all $0 \leq j \leq N$. However, this follows from the assumption that $\mathbf{c}(\mathbf{1}) = 1$, with a view on the explicit formula (4.4) for $F_{\mathcal{M}}$. Finally, $\text{Ch}(\mathcal{M})$ is even, that is, it vanishes on odd elements of $\mathbb{C}(\Omega)$. This follows directly from the fact that both \mathbf{c} and $\Phi_1^{\mathcal{M}}$ are parity-preserving (c.f. 4.4), since the supertrace vanishes on odd operators. The following was stated as Thm. B in the introduction.

Theorem 5.2 (Analyticity). *The Chern character $\text{Ch}(\mathcal{M})$ is an analytic cochain.*

Proof. That $\text{Ch}(\mathcal{M})$ satisfies the necessary estimates (3.6) follows directly from the fundamental estimate, Thm. 4.6. \square

We now proceed to show the closedness of $\text{Ch}(\mathcal{M})$, stated as Thm. A above.

Theorem 5.3 (Closedness). *The Chern character $\text{Ch}(\mathcal{M})$ is closed, that is,*

$$(\underline{d} + \underline{b} + \underline{B})\text{Ch}(\mathcal{M}) = 0.$$

A conceptual way to prove Thm. 5.3 is to exploit the coalgebra structure on the space $\text{Hom}(\mathbb{B}(\Omega), \mathcal{L}(\mathcal{H}))$ of $\mathcal{L}(\mathcal{H})$ -valued bar cochains, together with derived constructions. The result then follows from calculations similar to those of Quillen [47, Section 8]. However, instead of introducing the algebraic machinery to adapt Quillen's arguments, we use the trick of passing to the acyclic extension of \mathcal{M} as explained in Example 2.5.

Proof (of Thm. 5.3). Let us form the extended ϑ -summable Fredholm module $\mathcal{M}_{\mathbb{T}} := (\mathcal{H}, \mathbf{c}_{\mathbb{T}}, Q)$ over the dg algebra $\Omega_{\mathbb{T}} := \Omega[\sigma]$ with differential $d_{\mathbb{T}}$ (given by formula (2.7)), where σ is a formal variable of degree -1 with $\sigma^2 = 0$. Observe that it suffices to prove Thm. 5.3 for $\mathcal{M}_{\mathbb{T}}$ over $\Omega_{\mathbb{T}}$: Indeed, the inclusion $\Omega \subset \Omega_{\mathbb{T}}$ induces a map

$$j : \mathbb{C}(\Omega) \longrightarrow \mathbb{C}(\Omega_{\mathbb{T}}),$$

that commutes with all differentials, and one has $\text{Ch}(\mathcal{M}) = j^*\text{Ch}(\mathcal{M}_{\mathbb{T}})$. Hence the result for $\text{Ch}(\mathcal{M}_{\mathbb{T}})$ implies that for $\text{Ch}(\mathcal{M})$.

We note that $\Phi_1^{\mathcal{M}_{\mathbb{T}}}$ and $F_{\mathcal{M}_{\mathbb{T}}}$ descend to the quotient $\mathbf{B}(\underline{\Omega}_{\mathbb{T}})$; let us denote by $\underline{\Phi}_1^{\mathcal{M}_{\mathbb{T}}}$ the quotient map induced by $\Phi_1^{\mathcal{M}_{\mathbb{T}}}$. For the acyclic extension, the nonzero components of the curvature are given by

$$\begin{aligned} F_{\mathcal{M}_{\mathbb{T}}}^{(0)} &= Q^2, \\ F_{\mathcal{M}_{\mathbb{T}}}^{(1)}(\theta) &= \mathbf{c}(d\theta') - [Q, \mathbf{c}(\theta')] - \mathbf{c}(\theta'') \\ F_{\mathcal{M}_{\mathbb{T}}}^{(2)}(\theta_1, \theta_2) &= (-1)^{|\theta'_1|} (\mathbf{c}(\theta'_1\theta'_2) - \mathbf{c}(\theta'_1)\mathbf{c}(\theta'_2)), \end{aligned} \tag{5.1}$$

where again the elements of $\theta \in \Omega_{\mathbb{T}}$ have been written as $\theta = \theta' + \sigma\theta''$. Consider the parity-preserving map⁹

$$\alpha : \mathbf{C}(\Omega_{\mathbb{T}}) \longrightarrow \mathbf{B}(\underline{\Omega}_{\mathbb{T}}), \quad (\theta_0, \dots, \theta_N) \longmapsto \underline{\mathbf{N}}(\sigma\theta_0, \theta_1, \dots, \theta_N), \tag{5.2}$$

where

$$\underline{\mathbf{N}} : \mathbf{B}(\underline{\Omega}_{\mathbb{T}}) \longrightarrow \mathbf{B}(\underline{\Omega}_{\mathbb{T}})$$

is the quotient map of the *averaging operator*, given by

$$\mathbf{N} : \mathbf{B}(\Omega_{\mathbb{T}}) \longrightarrow \mathbf{B}(\Omega_{\mathbb{T}}), \quad (\theta_1, \dots, \theta_N) \longmapsto \sum_{k=1}^N (-1)^{n_k(n_N - n_k)} (\theta_{k+1}, \dots, \theta_N, \theta_1, \dots, \theta_k),$$

with $n_k = |\theta_1| + \dots + |\theta_k| - k$. By (5.1), we have $F_{\mathcal{M}_{\mathbb{T}}}(\sigma) = -1$; therefore an application of Lemma 5.4 below shows that (up to a sign) $\text{Ch}(\mathcal{M}_{\mathbb{T}})$ is the pullback of $\underline{\Phi}_1^{\mathbb{T}}$ under α , more precisely

$$\alpha^* \text{Str}(\underline{\Phi}_1^{\mathcal{M}_{\mathbb{T}}}) = \text{Str}(\underline{\Phi}_1^{\mathcal{M}_{\mathbb{T}}}\alpha) = -\text{Ch}(\mathcal{M}_{\mathbb{T}}). \tag{5.3}$$

Denote by $\mathbf{B}^{\natural}(\Omega_{\mathbb{T}}) \subset \mathbf{B}(\Omega_{\mathbb{T}})$ the image of \mathbf{N} , which is the space of cyclic bar chains. On this subspace, $\text{Str}(\Phi_1^{\mathbb{T}})$ is coclosed by Thm. 4.7, meaning that

$$\text{Str}\left(\Phi_1^{\mathcal{M}_{\mathbb{T}}}\left((d_{\mathbb{T}} + b')c\right)\right) = 0 \tag{5.4}$$

for all bar chains $c \in \mathbf{B}^{\natural}(\Omega_{\mathbb{T}})$. This follows from the fact that $\Phi_1((d_{\mathbb{T}} + b')\theta)$ is a sum of super-commutators of operators on \mathcal{H} in this case, as can be seen from the explicit formula (4.11).

We are therefore interested in the interaction of α with respect to the various differentials. Here straightforward calculations show that

$$\underline{b}'\alpha + \alpha\underline{b} = 0, \quad \underline{d}_{\mathbb{T}}\alpha + \alpha\underline{d}_{\mathbb{T}} = -h, \quad \alpha\underline{B} = \underline{S}h, \tag{5.5}$$

where \underline{b}' , $\underline{d}_{\mathbb{T}}$ are the differentials on the quotient $\mathbf{B}(\underline{\Omega}_{\mathbb{T}})$, h is given by

$$h : \mathbf{C}(\Omega_{\mathbb{T}}) \longrightarrow \mathbf{B}(\underline{\Omega}_{\mathbb{T}}), \quad (\theta_0, \dots, \theta_N) \longmapsto \underline{\mathbf{N}}(\theta_0, \theta_1, \dots, \theta_N),$$

⁹Here of course, $\mathbf{B}(\underline{\Omega}_{\mathbb{T}})$ denotes the space given by formula (3.1), but with the *space* $\underline{\Omega}_{\mathbb{T}}$ instead of the *algebra* $\Omega_{\mathbb{T}}$. Of course, the differentials d and b' descend to this quotient.

and

$$\underline{S} : \mathbf{B}(\underline{\Omega}_{\mathbb{T}}) \longrightarrow \mathbf{B}(\underline{\Omega}_{\mathbb{T}})$$

is the quotient map of the map S considered in (3.10). We obtain that

$$\alpha(\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B}) = (\underline{d}_{\mathbb{T}} + \underline{b}')\alpha + (\underline{S} - 1)h, \quad (5.6)$$

so with a view on (5.3), we have that

$$-(\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B})\text{Ch}(\mathcal{M}_{\mathbb{T}}) = \text{Str}\left(\Phi_1^{\mathcal{M}_{\mathbb{T}}}(\underline{d}_{\mathbb{T}} + \underline{b}')\alpha\right) + \text{Str}(\Phi_1^{\mathcal{M}_{\mathbb{T}}}(\underline{S} - 1)h) \quad (5.7)$$

The first summand on the right hand side is zero by (5.4), since α takes values in $\mathbf{B}^{\natural}(\underline{\Omega}_{\mathbb{T}})$. On the other hand, it follows from the special case (5.9) of Lemma 5.4 below that on the subspace $\mathbf{B}^{\natural}(\underline{\Omega}_{\mathbb{T}})$, we have

$$\begin{aligned} \text{Str}(\Phi_1^{\mathcal{M}_{\mathbb{T}}}S) &= - \sum_{N=0}^{\infty} (-1)^{N+1} \sum_{k=0}^N \text{Str}\left\{\underbrace{F_{\mathcal{M}_{\mathbb{T}}}^{\geq 1}, \dots, F_{\mathcal{M}_{\mathbb{T}}}^{\geq 1}}_k, 1, \underbrace{F_{\mathcal{M}_{\mathbb{T}}}^{\geq 1}, \dots, F_{\mathcal{M}_{\mathbb{T}}}^{\geq 1}}_{N-k}\right\}_Q \\ &= - \sum_{N=0}^{\infty} (-1)^{N+1} \text{Str}\left\{\underbrace{F_{\mathcal{M}_{\mathbb{T}}}^{\geq 1}, \dots, F_{\mathcal{M}_{\mathbb{T}}}^{\geq 1}}_N\right\}_Q = \text{Str}(\Phi_1^{\mathbb{T}}). \end{aligned} \quad (5.8)$$

Since h takes values in $\mathbf{B}^{\natural}(\underline{\Omega}_{\mathbb{T}})$, this shows that the second summand on the right hand side of (5.7) vanishes. This finishes the proof. \square

In the proof above, we used the following lemma.

Lemma 5.4. *For operators A_0, \dots, A_N satisfying the assumptions of Lemma 4.2, we have*

$$\sum_{j=0}^N (-1)^{k_j} \text{Str}(\{A_{j+1}, \dots, A_N, A_0, \dots, A_j\}_Q) = \text{Str}(A_0\{A_1, \dots, A_N\}_Q),$$

where $k_j = (|A_0| + \dots + |A_j|)(|A_{j+1}| + \dots + |A_N|)$.

Proof. First notice that by the cyclic permutation property of the trace, we have after a substitution in Δ_N that

$$\text{Str}(A_0\{A_1, \dots, A_N\}_Q) = (-1)^{k_j} \text{Str}(A_{j+1}\{A_{j+2}, \dots, A_N, A_0, \dots, A_j\}_Q)$$

for any $j = 0, \dots, N$. A special case of this is

$$\text{Str}(A_0\{A_1, \dots, A_j, 1, A_{j+1}, \dots, A_N\}_Q) = (-1)^{k_j} \text{Str}(\{A_{j+1}, \dots, A_N, A_0, \dots, A_j\}_Q).$$

On the other hand, the semigroup property and integration by parts yields

$$\{A_1, \dots, A_j, 1, A_{j+1}, \dots, A_N\}_Q = \int_{\Delta_N} (\tau_{j+1} - \tau_j) e^{-\tau_1 Q^2} \prod_{j=1}^N A_j e^{-(\tau_{j+1} - \tau_j) Q^2} d\tau$$

Therefore, using that $\sum_{j=0}^N(\tau_{j+1} - \tau_j) = 1$, we obtain

$$\sum_{j=0}^N \{A_1, \dots, A_j, 1, A_{j+1}, \dots, A_N\}_Q = \{A_1, \dots, A_N\}_Q. \quad (5.9)$$

Combining this with the observation before yields the claim. \square

We close this section by showing that the Chern character descends to the Chen normalized complexes in the case that \mathcal{M} is a ϑ -summable Fredholm module, not only a *weak* Fredholm module, i.e. when \mathcal{M} satisfies the identities (2.2).

Theorem 5.5 (Chen normalization). *The Chern character $\text{Ch}(\mathcal{M}_{\mathbb{T}})$ of a ϑ -summable Fredholm module is Chen normalized.*

Proof. We have to show that $\text{Ch}(\mathcal{M}_{\mathbb{T}})$ vanishes on $D^{\mathbb{T}}(\Omega)$. To begin with, the fact that $\text{Ch}(\mathcal{M}_{\mathbb{T}})$ vanishes on the image of $S - \text{id}$ and R is true also for weak Fredholm modules, as it follows from the calculation (5.8) above, respectively directly from the definition. It remains to show that if (2.2) holds, then for all $f \in \Omega^0$ and each $i \geq 1$, we have $\text{Ch}(\mathcal{M}_{\mathbb{T}})S_i^{(f)} = 0$ as well as $\text{Ch}(\mathcal{M}_{\mathbb{T}})[\underline{d} + \underline{b}, S_i^{(f)}] = 0$. The first identity follows directly from the definition of $\Phi_T^{\mathcal{M}_{\mathbb{T}}}$, since the assumptions (2.2) imply that

$$F(f) = 0 \quad \text{and} \quad F(f, \theta) = F(\theta, f) = 0$$

for any $\theta \in \Omega$. To see the second identity, observe first that

$$\begin{aligned} [(\underline{d} + \underline{b}), S_i^{(f)}](\theta_0, \dots, \theta_N) &= (\theta_0, \dots, \theta_i, df, \theta_{i+1}, \dots, \theta_N) \\ &\quad - (\theta_0, \dots, \theta_i f, \theta_{i+1}, \dots, \theta_N) \\ &\quad + (\theta_0, \dots, \theta_i, f\theta_{i+1}, \dots, \theta_N) \end{aligned} \quad (5.10)$$

for $0 \leq i \leq N - 1$, while for $i = N$, we have

$$[(\underline{d} + \underline{b}), S_N^{(f)}](\theta_0, \dots, \theta_N) = (\theta_0, \dots, \theta_N, df) - (\theta_0, \dots, \theta_N f) + (f\theta_0, \dots, \theta_N). \quad (5.11)$$

To compute $\Phi_1^{\mathcal{M}_{\mathbb{T}}}$ of this, we use the identities

$$\begin{aligned} F(df) &= -[Q^2, \mathbf{c}(f)], & F(f\theta_1, \theta_2) &= \mathbf{c}(f)F(\theta_1, \theta_2), \\ F(df, \theta) + F(f\theta) &= \mathbf{c}(f)F(\theta), & F(\theta_1 f, \theta_2) &= \mathbf{c}(f)F(\theta_1, \theta_2), \\ F(\theta, df) - F(\theta f) &= F(\theta)\mathbf{c}(f), & F(\theta_1 f, \theta_2) &= F(\theta_1, f\theta_2), \end{aligned}$$

which follow directly from (2.2). The theorem is then a consequence of the integration by parts lemma, Lemma 4.9, after a careful investigation of the terms appearing in the formula (4.9). \square

6 Homotopy Invariance of the Chern Character

In this section, we show that the Chern character defined above is invariant under suitable deformations of Fredholm modules. While the results of this section are not necessary for the remainder of the paper, we believe them to be of independent interest, in particular for the proof of the localization formula for the Chern character in the case of Example 2.3, see §9 below. Throughout, let Ω be a locally convex dg algebra.

Definition 6.1 (Homotopy of Fredholm modules). A *homotopy* of ϑ -summable (weak) Fredholm modules over Ω is a family $\mathcal{M}^s = (\mathcal{H}, \mathbf{c}^s, Q_s)$ of ϑ -summable (weak) Fredholm modules, $s \in [0,1]$, satisfying the following conditions.

(H1) The seminorms appearing in (A1), (A2) of Definition 2.1 can be chosen independently of s , and for all $T > 0$, we have

$$\sup_{s \in [0,1]} \mathrm{Tr}(e^{-TQ_s^2}) < \infty;$$

(H2) For all $s \in [0,1]$, the operators Q_s have the same domain of definition, and for each element $h \in \mathrm{dom}(Q_s)$, the map $s \mapsto Q_s h$ is a continuously differentiable curve in \mathcal{H} ; in particular, the derivative \dot{Q}_s is a densely defined operator on \mathcal{H} . Moreover, we require that $\dot{Q}_s \Delta_s^{-1/2}$ and $\Delta_s^{-1/2} \dot{Q}_s$ are bounded, where $\Delta_s = Q_s^2 + 1$, with uniform norm bound

$$\sup_{s \in [0,1]} \|\Delta_s^{-1/2} \dot{Q}_s\| + \sup_{s \in [0,1]} \|\dot{Q}_s \Delta_s^{-1/2}\| < \infty;$$

(H3) For all $\theta \in \Omega$, the map $s \mapsto \mathbf{c}^s(\theta)$ is continuously differentiable with respect to the strong operator topology.

The main result of this section is the following.

Theorem 6.2 (Homotopy invariance). *If $\mathcal{M}^s = (\mathcal{H}, \mathbf{c}^s, Q_s)$, $s \in [0,1]$, is an homotopy of ϑ -summable Fredholm modules, then the Chern characters $\mathrm{Ch}(\mathcal{M}_T^0)$ and $\mathrm{Ch}(\mathcal{M}_T^1)$ are homologous in the complex $\mathbf{N}^{\mathbb{T}, \alpha}(\Omega)$.*

We start with some auxiliary constructions. To this end, we fix a homotopy $\mathcal{M}^s = (\mathcal{H}, \mathbf{c}^s, Q_s)$, $s \in [0,1]$, of ϑ -summable (possibly weak) Fredholm modules and define a family of auxiliary $\mathcal{L}(\mathcal{H})$ -valued bar cochains

$$\Psi_T^{\mathcal{M}^s} := -T \int_0^1 \Phi_{uT}^{\mathcal{M}^s} \dot{\omega}_{\mathcal{M}^s} \Phi_{(1-u)T}^{\mathcal{M}^s} du. \quad (6.1)$$

In the following two propositions, we will abbreviate $\Psi_T^s := \Psi_T^{\mathcal{M}^s}$, $\Phi_T^s := \Phi_T^{\mathcal{M}^s}$ etc.

Proposition 6.3. *There exists a continuous seminorm ν on Ω such that for each $s \in [0,1]$, each $T > 0$, and all $\theta_1, \dots, \theta_N \in \Omega$, one has the estimate*

$$\|\Psi_T^s(\theta_1, \dots, \theta_N)\|_1 \leq e^{T/2} \mathrm{Tr}[e^{-TQ_s^2/2}] \frac{T^N}{\sqrt{N!}} \nu(\theta_1) \cdots \nu(\theta_N)$$

for the trace class norms. The same is true for $\Psi_T Q_s$ and $Q_s \Psi_T$ instead of Ψ_T .

Proof. Again, the argument for $\Psi_T^s Q_s$ and $Q_s \Psi_T^s$ is similar to the one for Ψ_T^s , so we restrict ourselves to the discussion of the latter case.

For suitable operators A_1, \dots, A_N, B on \mathcal{H} , we have the identity

$$\begin{aligned} & \{A_1, \dots, A_k, B, A_{k+1}, \dots, A_N\}_{TQ_s} \\ &= \int_0^1 u^k (1-u)^{N-k} \{A_1, \dots, A_k\}_{uTQ}^s B \{A_{k+1}, \dots, A_N\}_{(1-u)TQ_s} du. \end{aligned} \quad (6.2)$$

Using this and formula (4.9) for the quantization map, we obtain that $\Psi_T^s(\theta_1, \dots, \theta_N)$ is given by the formula

$$\begin{aligned} & \sum_{k=1}^N \sum_{M=1}^N (-T)^{M+1} \sum_{j=1}^M \left(\sum_{I \in \mathcal{P}_{M,N}} \{F_s(\theta_{I_1}), \dots, F_s(\theta_{I_{j-1}}), \dot{Q}_s, F_s(\theta_{I_j}), \dots, F_s(\theta_{I_M})\}_{TQ_s} \right. \\ & \quad \left. \sum_{\substack{I \in \mathcal{P}_{M,N} \\ I_j = \{k\}}} \{F_s(\theta_{I_1}), \dots, F_s(\theta_{I_{j-1}}), \dot{c}^s(\theta_k), F_s(\theta_{I_{j+1}}), \dots, F_s(\theta_{I_M})\}_{TQ_s} \right). \end{aligned}$$

Now each of the brackets can be estimated using Lemma 4.2, where to estimate the first bracket, we also use the bound on $\dot{Q}_s \Delta_s^{-1/2}$. We can then proceed as in the proof of Thm. 4.6. The uniformity in s follows from the assumptions on the admissible pair. \square

Proposition 6.4. *We have*

$$\frac{d}{ds} \Phi_T^s = \delta \Psi_T^s + [\omega_s, \Psi_T^s].$$

Proof (of Prop. 6.4). Similar to the proof of Lemma 2.2 (5) in [35], one proves the formula

$$\frac{d}{ds} \{A_1, \dots, A_N\}_{TQ_s} = -T \sum_{k=0}^N \{A_1, \dots, A_k, [Q_s, \dot{Q}_s], A_{k+1}, \dots, A_N\}_{TQ_s},$$

for operators A_1, \dots, A_N on \mathcal{H} , provided both sides are well-defined. Using this formula and an index shift, we obtain, similar to the proof of Thm. 4.7, that

$$\begin{aligned} \frac{d}{ds} \Phi_T^s &= \sum_{N=0}^{\infty} (-T)^{N+1} \sum_{k=1}^{N+1} \left\{ \underbrace{F_s^{\geq 1}, \dots, F_s^{\geq 1}}_k, [Q_s, \dot{Q}_s] + \dot{F}_s^{\geq 1}, \underbrace{F_s^{\geq 1}, \dots, F_s^{\geq 1}}_{N-k} \right\}_{TQ_s} \\ &= -T \int_0^1 \Phi_{uT}^s \dot{F}_s \Phi_{(1-u)T}^s du, \end{aligned}$$

where we used that $\dot{F}_s = [Q_s, \dot{Q}_s] + \dot{F}_s^{\geq 1}$, as well as the identity (6.2). Now

$$\delta \dot{\omega}_s + [\omega_s, \dot{\omega}_s] = \frac{d}{ds} \{\delta \omega_s + \omega_s^2\} = \dot{F}_s.$$

Therefore, using that $\delta\Phi_T^s + [\omega_s, \Phi_T^s] = 0$ by Thm. 4.7, we obtain

$$\int_0^1 \Phi_{sT}^s \dot{F}_s \Phi_{(1-s)T}^s ds = \delta \int_0^1 \Phi_{sT}^s \dot{\omega}_s \Phi_{(1-s)T}^s ds + \left[\omega_s, \int_0^1 \Phi_{sT}^s \dot{\omega}_s \Phi_{(1-s)T}^2 ds \right],$$

as requested. \square

In order to prove Thm. 6.2, fix a homotopy $\mathcal{M}^s := (\mathcal{H}, \mathbf{c}^s, Q_s)$, $s \in [0,1]$, of ϑ -summable Fredholm modules and let $\mathcal{M}_{\mathbb{T}}^s := (\mathcal{H}, \mathbf{c}_{\mathbb{T}}^s, Q_s)$, $s \in [0,1]$, be the associated homotopy of ϑ -summable *weak* Fredholm modules over $\Omega_{\mathbb{T}}$. Now using the map α defined in (5.2), we define the corresponding *Chern-Simons form* by

$$\text{CS}((\mathcal{M}_{\mathbb{T}}^s)_{s \in [0,1]}) := - \int_0^1 \alpha^* \text{Str}(\Psi_1^{\mathcal{M}_{\mathbb{T}}^s}) ds. \quad (6.3)$$

By Prop 6.3, $\text{CS}((\mathcal{M}_{\mathbb{T}}^s)_{s \in [0,1]})$ is an analytic cochain, and similar to the proof of Thm. 5.5, one shows that it is Chen normalized if \mathcal{M}^s satisfies (2.2). Thm. 6.2 is then a consequence of the following result.

Theorem 6.5 (Transgression formula). *For any $T > 0$, we have the transgression formula*

$$\text{Ch}(\mathcal{M}_{\mathbb{T}}^1) - \text{Ch}(\mathcal{M}_{\mathbb{T}}^0) = (\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B}) \text{CS}((\mathcal{M}_{\mathbb{T}}^s)_{s \in [0,1]}).$$

Proof. Since $-\text{Ch}(\mathcal{M}_{\mathbb{T}}^s) = \alpha^* \text{Str}(\Phi^{\mathcal{M}_{\mathbb{T}}^s}) = \text{Str}(\Phi^{\mathcal{M}_{\mathbb{T}}^s} \alpha)$ by the proof Thm. 5.3, Prop. 6.4 implies

$$-\frac{d}{ds} \text{Ch}(\mathcal{M}_{\mathbb{T}}^s) = \text{Str}(\delta \Psi_1^{\mathcal{M}_{\mathbb{T}}^s} \alpha) + \text{Str}([\omega^s, \Psi_1^{\mathcal{M}_{\mathbb{T}}^s}] \alpha) = \text{Str}(\Psi_1^{\mathcal{M}_{\mathbb{T}}^s} (d_{\mathbb{T}} + b') \alpha),$$

where we used that α takes values in the space $\mathbf{B}^{\natural}(\underline{\Omega}_{\mathbb{T}})$ of cyclic chains, on which $[\omega^s, \Psi_1^{\mathcal{M}_{\mathbb{T}}^s}]$ vanishes. Using (5.6), we then get

$$\frac{d}{ds} \text{Ch}(\mathcal{M}_{\mathbb{T}}^s) = -(\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B}) \alpha^* \text{Str}(\Psi_1^{\mathcal{M}_{\mathbb{T}}^s}) - \text{Str}(\Psi_1^{\mathcal{M}_{\mathbb{T}}^s} (S - \text{id}) h).$$

The second summand on the right hand side vanishes by an argument similar to that in the proof Thm. 5.3, which finishes the proof. \square

Remark 6.6. Of course, similar results hold in the case of a homotopy \mathcal{M}^s , $s \in [0,1]$, of ϑ -summable *weak* Fredholm modules over Ω . In this case $\text{Ch}(\mathcal{M}_{\mathbb{T}}^0)$ and $\text{Ch}(\mathcal{M}_{\mathbb{T}}^1)$ are homologous in the complex $\mathbf{C}_{\alpha}(\Omega_{\mathbb{T}})$.

7 The Bismut-Chern Character of an Idempotent

Let Ω be a locally convex dg algebra. The algebra of $n \times n$ matrices with entries in Ω is denoted by $\text{Mat}_n(\Omega)$, which is then again a locally convex dg algebra, with the differential (also denoted by d) acting entrywise. We can form the acyclic extension $\Omega_{\mathbb{T}}$ of Ω as in Example 2.5 by adjoining a formal variable σ of degree -1 . The same can be done for $\text{Mat}_n(\Omega)$; we have $\text{Mat}_n(\Omega)_{\mathbb{T}} \cong \text{Mat}_n(\Omega_{\mathbb{T}})$.

The goal of this section is to construct the Bismut-Chern character associated to an idempotent p in the algebra $\text{Mat}_n(\Omega^0)$, which will be an even element of $\mathbb{C}^\epsilon(\Omega_{\mathbb{T}})$, closed as an element in the complex $\mathbb{N}^{\mathbb{T},\epsilon}(\Omega)$. To this end, let $p \in \text{Mat}_n(\Omega^0)$ be an idempotent, $p^2 = p$, and write $p^\perp := \mathbf{1} - p$. The formula

$$D\theta = p d(p\theta) + p^\perp d(p^\perp\theta). \quad (7.1)$$

then defines an operator D on $\text{Mat}_n(\Omega)$. The corresponding *connection form* $\varrho_p \in \text{Mat}_n(\Omega^1)$ is defined by the equation $D = d + \varrho_p$. Its *curvature* is the element R_p of $\text{Mat}_n(\Omega^2)$ given by $R_p = d\varrho_p + \varrho_p^2$. We have the explicit formulas

$$\varrho_p = p dp + p^\perp dp^\perp = (2p - \mathbf{1})dp, \quad R_p = (dp)^2.$$

for the connection form and the curvature.

Example 7.1 (Vector bundles with connection). Given a manifold X , any smooth function $p : X \rightarrow \text{Mat}_n(\mathbb{C})$ taking values in idempotents (equivalently, an idempotent in $\text{Mat}_n(C^\infty(X)) \subset \text{Mat}_n(\Omega(X))$) determines a vector bundle $E := \text{im}(p)$ with connection D as defined above. It is well-known [45, Thm. 1] that any vector bundle with connection on X is isomorphic (through a connection-preserving isomorphism) to one of this form.

Following [34, Section 6], we combine these elements into a single even element $\mathfrak{R}_p \in \text{Mat}_n(\Omega_{\mathbb{T}})$ given by

$$\mathfrak{R}_p := \varrho_p + \sigma R_p.$$

Definition 7.2 (Bismut-Chern character). The *Bismut-Chern character* of p is the even entire chain $\text{Ch}(p) \in \mathbb{C}_+^\epsilon(\Omega_{\mathbb{T}})$ given by

$$\text{Ch}(p) = \sum_{N=0}^{\infty} (-1)^N \text{tr}(p, \underbrace{\mathfrak{R}_p, \dots, \mathfrak{R}_p}_N).$$

In the definition,

$$\text{tr} : \mathbb{C}^\epsilon(\text{Mat}_n(\Omega_{\mathbb{T}})) \longrightarrow \mathbb{C}^\epsilon(\Omega_{\mathbb{T}}), \quad (\Theta_0, \dots, \Theta_N) \longmapsto \sum_{i_0, \dots, i_N=1}^n ((\Theta_0)_{i_0}^{i_1}, (\Theta_1)_{i_1}^{i_2}, \dots, (\Theta_N)_{i_N}^{i_0})$$

is the *generalized trace map*, which preserves all differentials, see [42, 1.2.1]. Note that $\text{Ch}(p)$ is indeed entire: If ν is a continuous seminorm on $\Omega_{\mathbb{T}}$, then

$$\pi_{\nu, N}(p, \underbrace{\mathfrak{R}_p, \dots, \mathfrak{R}_p}_N) \leq \nu(p)\nu(\mathfrak{R}_p)^N,$$

so that clearly $\epsilon_\nu(\text{Ch}(p)) < \infty$. Moreover, $\text{Ch}(p)$ is even because \mathfrak{R}_p is odd in $\text{Mat}_n(\Omega_{\mathbb{T}})$, hence even in $\text{Mat}_n(\Omega_{\mathbb{T}})[1]$.

Remark 7.3. The entire chain $\text{Ch}(p)$ was first considered by Getzler, Jones and Petrack in [34, Section 6], in the Example 2.3, where $\Omega = \Omega(X)$, differential forms on a manifold X . The significance of $\text{Ch}(p)$ is that under the extended or equivariant iterated integral map (cf. §9 below), $\text{Ch}(p)$ is sent to the *Bismut-Chern character* $\text{BCh}(E, \nabla)$ of the vector bundle with connection (E, ∇) corresponding to p as in Example 7.1. This is an equivariantly closed differential form on the loop space, first considered by Bismut in [10]. An odd variant of $\text{Ch}(p)$ has been recently constructed by the first named author and S. Cacciatori in [17], producing the odd Bismut-Chern character of Wilson [51].

Theorem 7.4. *As an element of the Chen normalized complex $\mathbf{N}^{\mathbb{T}, \epsilon}(\Omega)$, one has*

$$(\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B})\text{Ch}(p) = 0.$$

For the proof of the proof of Thm. 7.4, we need the following Lemma; throughout, we write \mathfrak{R} instead of \mathfrak{R}_p .

Lemma 7.5. *We have the identities*

$$d_{\mathbb{T}}p + [\mathfrak{R}, p] = 0 \tag{7.2}$$

$$d_{\mathbb{T}}\mathfrak{R} + \mathfrak{R}^2 = 0 \tag{7.3}$$

$$d_{\mathbb{T}}(\sigma p) + [\mathfrak{R}, \sigma p] = -p. \tag{7.4}$$

Proof. This follow from straightforward calculation, using the relations

$$[p, \varrho] = dp, \quad [\varrho, dp] = [dp, p], \quad \text{and} \quad [R, p] = 0.$$

The first of these latter identities follows from differentiating the equation $p^2 = p$ and straightforward calculation using the definition of ϱ , the second identity follows from differentiating the first, and the third is a direct calculation using the previous results. \square

Remark 7.6. In fact, the proof shows that the element defined above is even closed *before* taking the trace, i.e. as an element in $\mathbf{C}^\epsilon(\text{Mat}_n(\Omega_{\mathbb{T}}))$.

Proof (of Thm. 7.4). We have to show that $(\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B})\text{Ch}(p)$ is contained in the closure of $\mathbf{D}^{\mathbb{T}}(\Omega)$ in $\mathbf{C}^{\mathbb{T}, \epsilon}(\Omega)$. To this end, calculate

$$\begin{aligned} \underline{d}_{\mathbb{T}}(p, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_N) &= (d_{\mathbb{T}}p, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_N) - \sum_{k=1}^N (p, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_{k-1}, d_{\mathbb{T}}\mathfrak{R}, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_{N-k}) \\ \underline{b}(p, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_N) &= ([p, \mathfrak{R}], \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_{N-1}) + \sum_{k=1}^{N-1} (p, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_{k-1}, \mathfrak{R}^2, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_{N-k-1}). \end{aligned}$$

By Lemma 7.5, the second equation can be rewritten as

$$\underline{b}(p, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_N) = (d_{\mathbb{T}}p, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_{N-1}) - \sum_{k=1}^{N-1} (p, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_{k-1}, d_{\mathbb{T}}\mathfrak{R}, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_{N-k-1}),$$

which telescopes with the the right hand side of the first equation as we take the alternating sum over N , meaning that

$$(\underline{d}_{\mathbb{T}} + \underline{b})\text{Ch}(p) = 0.$$

Now $\underline{B}\text{Ch}(p)$ is a sum of the terms $\underline{B}(p, \mathfrak{R}, \dots, \mathfrak{R})$, which are contained in $D^{\mathbb{T}}(\Omega)$, since p is of degree zero. \square

We proceed by discussing the homotopy invariance of the Bismut-Chern character. First some notation: If $(\theta_j^s)_{s \in [0,1]}$, $j = 0, \dots, N$ are continuously differentiable families of elements in Ω , we set

$$\partial_s(\theta_0^s, \dots, \theta_N^s) := - \sum_{k=0}^N (-1)^{m_{k-1}} (\theta_0^s, \dots, \theta_{k-1}^s, \sigma \frac{d}{ds} \theta_k^s, \theta_{k+1}^s, \dots, \theta_N^s),$$

where $m_k = |\theta_0| + \dots + |\theta_k| - k$ and θ_k^s denotes the derivative in direction of s . Now given a continuously differentiable family $(p_s)_{s \in [0,1]}$ of idempotents, we define the corresponding *Bismut-Chern-Simons form* by

$$\text{CS}((p_s)_{s \in [0,1]}) := \int_0^1 \partial_s \text{Ch}(p_s) ds.$$

As for $\text{Ch}(p)$, it can be easily shown that $\text{Ch}(p_s)$ is a continuously differentiable function of s with values in $C^\epsilon(\Omega_{\mathbb{T}})$ and that $\partial_s \text{Ch}(p_s)$ is entire for any $s \in [0, 1]$. Now since this is a continuous function of s with values in the complete locally convex space $C^\epsilon(\Omega_{\mathbb{T}})$, its integral $\text{CS}((p_s)_{s \in [0,1]})$ is well-defined.

Proposition 7.7 (Transgression). *As elements of the Chen normalized complex $N^{\mathbb{T}, \epsilon}(\Omega)$, we have the transgression formula*

$$\text{Ch}(p_1) - \text{Ch}(p_0) = (\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B})\text{CS}((p_s)_{s \in [0,1]}).$$

Proof. We have to show that $(\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B})\text{CS}((p_s)_{s \in [0,1]})$ is contained in the closure $\overline{D^{\mathbb{T}}(\Omega)}$ of $D^{\mathbb{T}}(\Omega)$ in $C^\epsilon(\Omega_{\mathbb{T}})$, for all $s \in [0,1]$. To this end, it is straightforward to verify that

$$\underline{d}\partial_s + \partial_s \underline{d} = \underline{b}\partial_s + \partial_s \underline{b} = \underline{B}\partial_s + \partial_s \underline{B} = 0, \quad \text{and} \quad \underline{l}\partial_s + \partial_s \underline{l} = \frac{d}{ds},$$

where

$$\frac{d}{ds}(\theta_0^s, \dots, \theta_N^s) = \sum_{k=0}^N (\theta_0^s, \dots, \frac{d}{ds} \theta_k^s, \dots, \theta_N^s)$$

is the derivative in the locally convex space $C^\epsilon(\Omega_{\mathbb{T}})$. Hence we have

$$\begin{aligned} (\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B})\text{CS}((p_s)_{s \in [0,1]}) &= - \int_0^1 \partial_s(\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B})\text{Ch}(p_s)ds + \int_0^1 \frac{d}{ds}\text{Ch}(p_s)ds \\ &= - \int_0^1 \partial_s \underline{B}\text{Ch}(p_s)ds + \text{Ch}(p_1) - \text{Ch}(p_0), \end{aligned}$$

where in the last step, we used the fundamental theorem of calculus and the fact that $(\underline{d}_{\mathbb{T}} + \underline{b})\text{Ch}(p_s) = 0$ by the proof of Thm. 7.4 above.

It therefore remains to show that $\partial_s \underline{B}\text{Ch}(p_s)$ is contained in $\overline{\mathbb{D}^{\mathbb{T}}(\Omega)}$ for all s , which is not quite obvious as ∂_s does not generally preserve $\overline{\mathbb{D}^{\mathbb{T}}(\Omega)}$. It is easy to see that $\partial_s \underline{B}\text{Ch}(p_s)$ is entire, so we just need an algebraic argument. To this end, observe that modulo $\overline{\mathbb{D}^{\mathbb{T}}(\Omega)}$,

$$\begin{aligned} \underline{B}\text{Ch}(p_s) &= \sum_{N=0}^{\infty} (-1)^N \sum_{k=0}^N (\mathbf{1}, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_k, \sigma \dot{p}, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_{N-k}) \\ &= \underline{B} \sum_{N=0}^{\infty} (-1)^N (\sigma \dot{p}, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_N), \end{aligned} \tag{7.5}$$

where we dropped the dependence on s for readability and wrote $\dot{p} = \frac{d}{ds}p_s$. Now because of $\dot{p} = \dot{p}p + p\dot{p}$ and the identity (5.10), we have modulo $\overline{\mathbb{D}^{\mathbb{T}}(\Omega)}$

$$(\sigma \dot{p}, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_N) = (p\sigma \dot{p}, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_N) + (\sigma \dot{p}, \underbrace{d_{\mathbb{T}}p, \mathfrak{R}, \dots, \mathfrak{R}}_N) + (\sigma \dot{p}, \underbrace{p\mathfrak{R}, \mathfrak{R}, \dots, \mathfrak{R}}_{N-1}).$$

After summing over N and shifting an index on the second factor, we can use (7.2) to obtain that modulo $\overline{\mathbb{D}^{\mathbb{T}}(\Omega)}$,

$$\sum_{N=0}^{\infty} (-1)^N (\sigma \dot{p}, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_N) = \sum_{N=0}^{\infty} (-1)^N (p\sigma \dot{p}, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_N) + \sum_{N=0}^{\infty} (-1)^N (\sigma \dot{p}, \underbrace{\mathfrak{R}p, \mathfrak{R}, \dots, \mathfrak{R}}_{N-1}).$$

Repeating this, we can shift the factor of p in the second sum all the way to the right and then use (7.2) combined with (5.11), making this term cancel with the first sum on the right hand side of the above equation. This implies

$$\sum_{N=0}^{\infty} (-1)^N (\sigma \dot{p}, \underbrace{\mathfrak{R}, \dots, \mathfrak{R}}_N) \in \overline{\mathbb{D}^{\mathbb{T}}(\Omega)}.$$

The claim now follows from (7.5), bearing in mind that \underline{B} preserves $\overline{\mathbb{D}^{\mathbb{T}}(\Omega)}$. \square

It is easy to see that $\text{Ch}(p \oplus q) = \text{Ch}(p) + \text{Ch}(q)$, for projections $p \in \text{Mat}_n(\Omega^0)$, $q \in \text{Mat}_m(\Omega^0)$, where $p \oplus q \in \text{Mat}_{n+m}(\Omega^0)$. The transgression formula above therefore implies the following result.

Theorem 7.8. *The Bismut-Chern character defines a group homomorphism*

$$K_0(\Omega^0) \longrightarrow H^+(\mathbb{N}^{\mathbb{T}, \epsilon}(\Omega))$$

from the K -theory of the algebra Ω to the cohomology of the complex $\mathbb{N}^{\mathbb{T}, \epsilon}(\Omega)$.

8 An Index Theorem

In this section, let Ω be a locally convex dg algebra and let $\mathcal{M} = (\mathcal{H}, \mathbf{c}, Q)$ be a ϑ -summable Fredholm module over Ω . We emphasize that to obtain the results of this section, it is crucial that \mathcal{M} satisfies (2.2).

Given an idempotent $p = p^2 \in \text{Mat}_n(\Omega^0)$, (2.2) implies that the operator $\mathbf{c}(p)$ is an even idempotent acting on the Hilbert space \mathcal{H}^n , so that $\mathbf{c}(p)\mathcal{H}^n$ becomes a closed subspace of \mathcal{H} and thus a Hilbert space on its own right. As $\mathbf{c}(p)$ is even, the \mathbb{Z}_2 -grading restricts to $\mathbf{c}(p)\mathcal{H}^n$. We obtain a well-defined odd unbounded operator

$$Q_p := \mathbf{c}(p)Q\mathbf{c}(p),$$

which is densely defined and closed on the Hilbert space $\mathbf{c}(p)\mathcal{H}^n$. The purpose of this section is to prove the following index theorem.

Theorem 8.1 (Geometric index theorem). *The operator Q_p defined above is Fredholm and its \mathbb{Z}_2 -graded index is given by the formula*

$$\text{ind}(Q_p) = \langle \text{Ch}(\mathcal{M}_{\mathbb{T}}), \text{Ch}(p) \rangle. \quad (8.1)$$

This theorem can be seen as a “loop space version” of the index theorem for the JLO-cocycle of Getzler-Szenes [35]. While their proof relies on the homotopy invariance of the Chern character, a remarkable observation is that in our setup, the right hand side of (8.1) is *directly* the super trace of a heat operator, without having to defer to any homotopy result. Precisely, we have the following proposition.

Proposition 8.2 (Perturbation series). *Let $p \in \text{Mat}_n(\Omega^0)$ be an idempotent. Then the unbounded operator*

$$Q_1 := \mathbf{c}(p)Q\mathbf{c}(p) + \mathbf{c}(p^\perp)Q\mathbf{c}(p^\perp),$$

on \mathcal{H}^n is closed on the same domain as Q , and its square generates a strongly continuous semigroup of operators. Moreover, for all $T > 0$, one has the formula

$$e^{-TQ_1^2} = \sum_{N=0}^{\infty} (-1)^N \Phi_T^{\mathcal{M}_{\mathbb{T}}}(\underbrace{\mathfrak{R}_p, \dots, \mathfrak{R}_p}_N). \quad (8.2)$$

Proof. From (2.2), we obtain $[Q, \mathbf{c}(p)] = \mathbf{c}(dp)$ and $\mathbf{c}(p)\mathbf{c}(dp) = \mathbf{c}(p dp)$. Therefore

$$Q_1 = Q + \mathbf{c}(2p - \mathbf{1})[Q, \mathbf{c}(p)] = Q + \mathbf{c}((2p - \mathbf{1})dp).$$

This shows that that Q_1 is in fact a bounded perturbation of Q , hence is a closed operator on the same domain as Q . On the other hand, we have $p dp + p^\perp dp^\perp = (2p - \mathbf{1})dp$, so with a view on (5.1),

$$\begin{aligned} F(\mathfrak{R}) &= \mathbf{c}(d\{(2p - \mathbf{1})dp\}) - [Q, \mathbf{c}((2p - \mathbf{1})dp)] - \mathbf{c}((dp)^2) \\ &= \mathbf{c}((dp)^2) - [Q, \mathbf{c}((2p - \mathbf{1})dp)], \end{aligned}$$

as well as

$$\begin{aligned} F(\mathfrak{A}, \mathfrak{A}) &= \mathbf{c}((2p-1)dp)^2 - \mathbf{c}(((2p-1)dp)^2) \\ &= \mathbf{c}((2p-1)dp)^2 + \mathbf{c}((dp)^2). \end{aligned}$$

Putting together, we obtain

$$Q^2 - F(\mathfrak{A}) + F(\mathfrak{A}, \mathfrak{A}) = Q^2 + [Q, \mathbf{c}((2p-1)dp)] + \mathbf{c}((2p-1)dp) = Q_1^2.$$

Hence we can write $Q_1^2 = Q^2 + R$, where $R := F(\mathfrak{A}, \mathfrak{A}) - F(\mathfrak{A})$. By the assumptions (A1), (A2) on the Fredholm module, we can use Lemma 4.2 to define $e^{-TQ_1^2} = e^{-T(Q^2+R)}$ using the perturbation series

$$e^{-T(Q^2+R)} = \sum_{M=0}^{\infty} (-T)^M \underbrace{\{R, \dots, R\}}_M{}_{TQ}. \quad (8.3)$$

It is easy to see that this indeed defines a strongly continuous semigroup of operators with infinitesimal generator $Q^2 + P = Q_1^2$. Now we have

$$\sum_{M=0}^{\infty} (-T)^M \underbrace{\{R, \dots, R\}}_M{}_{TQ} = \sum_{M=0}^{\infty} (-T)^M \{F(\mathfrak{A}, \mathfrak{A}) - F(\mathfrak{A}), \dots, F(\mathfrak{A}, \mathfrak{A}) - F(\mathfrak{A})\}_{TQ}.$$

Expanding each term on the right hand side of this equation by multi-linearity, we see that we obtain an infinite sum of brackets $\{\dots\}_{TQ}$ which contains all possible sequences of operators $F(\mathfrak{A})$ and $F(\mathfrak{A}, \mathfrak{A})$. Observe now that by formula (4.9), this coincides with the right hand side of (8.2). \square

Proof (of Thm. 8.1). If $\mathbf{c}(p)$ is self-adjoint, the proof follows directly from Prop. 8.2. Namely, in that case, we get

$$\langle \text{Ch}(\mathcal{M}_{\mathbb{T}}), \text{Ch}(p) \rangle = \text{Str}_{\mathcal{H}^n}(\mathbf{c}(p)e^{-Q_1^2}) = \text{Str}_{\mathbf{c}(p)\mathcal{H}^n}(e^{-Q_p^2}),$$

which equals the graded index of Q_p by the usual McKean-Singer formula.

To deal with the general case, we adapt an idea of Getzler-Szenes [35] to our setup, making the following construction. Consider the $*$ -subalgebra

$$\mathcal{B}_Q := \left\{ A \in \mathcal{L}(\mathcal{H})^+ \mid \Delta^{1/2}A\Delta^{-1/2} \text{ and } \Delta^{1/2}A^*\Delta^{-1/2} \right. \\ \left. \text{are densely defined and bounded} \right\}, \quad (8.4)$$

of $\mathcal{L}(\mathcal{H})$, where $\Delta = Q^2 + 1$ and $\mathcal{L}(\mathcal{H})^+$ denotes the subalgebra of even operators. Note that we have

$$(Q + A)^2 = Q^2 + [Q, A] + A^2,$$

hence for $A \in \mathcal{B}_Q$, the operator $R = [Q, A] + A^2$ satisfies the assumptions of Lemma 4.2. Hence for any $A \in \mathcal{B}_Q$, the heat semigroup $e^{-T(Q+A)^2}$ can be defined using the perturbation series (8.3).

Furthermore, set

$$\mathcal{A}_Q := \{A \in \mathcal{L}(\mathcal{H}) \mid [Q, A] \in \mathcal{B}_Q\}$$

and let $\Omega_Q = \Omega_{\mathcal{A}_Q}$ the corresponding dg algebra of non-commutative differential forms, which comes with a canonical map $\mathbf{c}_Q : \Omega_Q \rightarrow \mathcal{L}(\mathcal{H})$ so that $\mathcal{M}_Q = (\mathcal{H}, \mathbf{c}_Q, Q)$ is a Fredholm module over Ω_Q , as explained in Example 2.2.

Now the calculations in the proof of Prop. 8.2 show that for an idempotent $p \in \Omega^0$, we have

$$\langle \text{Ch}(\mathcal{M}_{\mathbb{T}}), \text{Ch}(p) \rangle = \langle \text{Ch}(\mathcal{M}_{Q, \mathbb{T}}), \text{Ch}(\mathbf{c}(p)) \rangle,$$

where the right hand side denotes the pairing of the extended entire cyclic complex associated to Ω_Q . Therefore, it suffices to prove Thm. 8.1 for the “universal example” Ω_Q .

To this end, we will show that the idempotent $\mathbf{c}(p)$ can be homotoped within \mathcal{A}_Q to a self-adjoint projection P , as then

$$\langle \text{Ch}(\mathcal{M}_{Q, \mathbb{T}}), \text{Ch}(\mathbf{c}(p)) \rangle = \langle \text{Ch}(\mathcal{M}_{Q, \mathbb{T}}), \text{Ch}(P) \rangle$$

by the transgression formula Thm. 7.7 and the closedness of $\text{Ch}(\mathcal{M}_{Q, \mathbb{T}})$, Thm. 5.3 (together with the fact that it is Chen normalized, Thm. 5.5). By Prop. 4.6.2 in [15], this property follows from the fact that \mathcal{A}_Q is a *spectral* subalgebra of $\mathcal{L}(\mathcal{H})$, i.e. that for any $A \in \mathcal{A}_Q$ which is invertible in $\mathcal{L}(\mathcal{H})$, its inverse $A^{-1} \in \mathcal{A}_Q$. To see that the larger algebra \mathcal{B}_Q is spectral, notice that for $A \in \mathcal{B}_Q$,

$$\Delta^{1/2} A^{-1} \Delta^{-1/2} = (\Delta^{1/2} A \Delta^{-1/2})^{-1}.$$

But $\Delta^{1/2} A \Delta^{-1/2}$ is bounded, hence its inverse is bounded as well. Now for $A \in \mathcal{A}_Q$, we have

$$[Q, A^{-1}] = A^{-1}[Q, A]A^{-1},$$

which is a product of elements in \mathcal{B}_Q by assumption, hence in \mathcal{B}_Q . Since these arguments also work for A^* instead of A , this shows that \mathcal{A}_Q is indeed spectral in $\mathcal{L}(\mathcal{H})$ and finishes the proof. \square

Thm. 8.1 in combination with Thm. 7.7 and homotopy invariance of the index easily implies the following result.

Corollary 8.3 (Homological index theorem). *We have a commutative diagram*

$$\begin{array}{ccc} K_0(\Omega^0) & \xrightarrow{\text{Ch}} & H^+(\mathbf{N}^{\mathbb{T}, \epsilon}(\Omega)) \\ \text{ind}_{\mathcal{M}} \downarrow & & \downarrow \text{Ch}(\mathcal{M}_{\mathbb{T}}) \\ \mathbb{Z} & \longrightarrow & \mathbb{C}, \end{array} \tag{8.5}$$

where $\text{ind}_{\mathcal{M}}$ is the map that assigns the index of the operator Q_p on $\mathbf{c}(p)\mathcal{H}$.

Notice in particular that if \mathcal{A} is a locally convex algebra, we have such a commutative diagram (8.5) for any dg algebra Ω with $\Omega^0 = \mathcal{A}$ together with a ϑ -summable Fredholm module \mathcal{M} over Ω .

9 The Supersymmetric Path Integral and its Localization Property

In this section, we finally discuss the application of our abstract theory to the construction of the supersymmetric path integral for the $N = 1/2$ supersymmetric σ -model and its localization property, which originally was our main motivation.

Let X a compact spin manifold of even dimension n . In the considerations of Atiyah [5], the corresponding supersymmetric path integral is an integration functional I for suitable differential forms on the smooth loop space LX of X , formally defined by (1.1). The aim of this section is to use the abstract results from the previous sections to rigorously construct this linear functional and to establish its two fundamental properties: supersymmetry and its localization property (1.2).

To this end, consider the dg algebra $\Omega(X)$ of differential forms over X . There is a canonical Fredholm module $\mathcal{M}^X = (\mathcal{H}, Q, \mathbf{c})$ over $\Omega(X)$, as discussed in Example 2.3: $\mathcal{H} = L^2(X, \Sigma)$, the space of square-integrable sections of the spinor bundle Σ , $Q = \mathbb{D}$, the Dirac operator, and \mathbf{c} is given by Clifford multiplication (2.6). This Fredholm module satisfies the identities (2.2), hence according to Thm. 5.5, the Chern character $\text{Ch}(\mathcal{M}_{\mathbb{T}}^X)$ is Chen normalized, i.e. descends to a cochain on the complex $\mathbf{N}^{\epsilon, \mathbb{T}}(\Omega(X))$ and itself defines an element in $\mathbf{N}_{\mathbb{T}, \alpha}^+(\Omega(X))$. The following result is key in this situation; it seems likely that it can be generalized to more general situations analogously to the local index formula of Connes and Moscovici, c.f. [24, 38].

Theorem 9.1 (Localization). *As an element of the complex $\mathbf{N}_{\mathbb{T}, \alpha}^+(\Omega(X))$, the Chern character $\text{Ch}(\mathcal{M}_{\mathbb{T}}^X)$ is cohomologous to the cochain μ_0 defined by*

$$\mu_0(\theta_0, \dots, \theta_N) = \frac{1}{(2\pi i)^{n/2} N!} \int_X \widehat{A}(X) \wedge \theta'_0 \wedge \theta''_1 \wedge \dots \wedge \theta''_N \quad (9.1)$$

for $\theta_0, \dots, \theta_N \in \Omega(X)_{\mathbb{T}}$. Here

$$\widehat{A}(X) = \det^{1/2} \left(\frac{R/2}{\sinh(R/2)} \right) \quad (9.2)$$

is the Chern-Weil representative of the \widehat{A} -form of X , involving the Riemann curvature tensor R of X (c.f. [7, (1.35)]).

Direct computations show that $\underline{d}\mu_0 = 0$ (this uses that $\widehat{A}(X)$ is closed together with Stokes theorem), $\underline{b}\mu_0 = 0$ and $\underline{B}\mu_0 = \underline{\nu}\mu_0$, so that $(\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B})\mu_0 = 0$. Moreover, it is easy to check that μ_0 is both analytic and Chen normalized, hence indeed is a closed element of the complex $\mathbf{N}_{\mathbb{T}, \alpha}^+(\Omega(X))$.

Proof. We define a homotopy of ϑ -summable Fredholm modules $\mathcal{M}^{X, T} = (\mathcal{H}, Q_T, \mathbf{c}_T)$ over $\Omega(X)$ by setting

$$Q_T = T^{1/2} \mathbb{D}, \quad \mathbf{c}_T(\theta) = T^{|\theta|/2} \mathbf{c}(\theta).$$

One checks that the scaling is such that the relations (2.2) are preserved, hence we obtain Chen normalized Chern characters $\text{Ch}(\mathcal{M}_{\mathbb{T}}^{X,T}) \in \mathbf{N}_{\mathbb{T},\alpha}(\Omega(X))$, which are all cohomologous by Thm. 6.2. The theorem now follows from the following proposition, which generalizes the result of Block and Fox [16, Thm. 4.1] on the JLO-cocycle to our situation; c.f. also the calculations in [23]. \square

Proposition 9.2. *For all $\theta_0, \dots, \theta_N \in \Omega(X)_{\mathbb{T}}$, we have*

$$\lim_{T \rightarrow 0} \langle \text{Ch}(\mathcal{M}_{\mathbb{T}}^{X,T}), (\theta_0, \dots, \theta_N) \rangle = \mu_0(\theta_0, \dots, \theta_N).$$

Proof. Explicit calculations show that if $\theta_0, \dots, \theta_N \in \Omega(X)_{\mathbb{T}}$ are homogeneous (with, say $|\theta_k| = \ell_k$), we have

$$\begin{aligned} \langle \text{Ch}(\mathcal{M}_{\mathbb{T}}^{X,T}), (\theta_0, \dots, \theta_N) \rangle &= T^{m_N/2} \text{Str}(\mathbf{c}(\theta'_0) \Phi_T^{\mathcal{M}_{\mathbb{T}}^X}(\theta_1, \dots, \theta_N)) \\ &= \sum_{M=1}^N (-T)^M \sum_{\mathcal{P}_{M,N}} \int_{\Delta_M} \text{Str}(\mathbf{c}(\theta'_0) e^{-T\tau_1 \mathbb{D}^2} F(\theta_{I_1}) e^{-T(\tau_2 - \tau_1) \mathbb{D}^2} \dots F(\theta_{I_M}) e^{-T(1 - \tau_M) \mathbb{D}^2}) d\tau \end{aligned} \quad (9.3)$$

where $m_N = \ell_0 + \dots + \ell_N - N$. The right hand side is now amenable to Getzler rescaling [29, 7]; we assume for the rest of the proof that the reader is familiar with this technique. To calculate the short-time limit of the right hand side of (9.3), we need to calculate the orders of the operators $F(\theta_k)$, $F(\theta_k, \theta_{k+1})$ in the Getzler filtration. As usual, write $\theta = \theta' + \sigma\theta''$ so that $|\theta'_k| = \ell_k$, $|\theta''_k| = \ell_k + 1$. Since Clifford multiplication with a differential form θ has order $|\theta|$ in the Clifford filtration and the Dirac operator has degree two, the operator

$$F(\theta_k) = \mathbf{c}(d\theta'_k) - [\mathbb{D}, \mathbf{c}(\theta'_k)] - \mathbf{c}(\theta''_k) \quad (9.4)$$

has degree $\ell_k + 1$. However, an explicit calculation of the commutator (extending the formula of Prop. 3.45 of [7] to arbitrary form degrees) shows that the terms involving θ'_k are actually of order ℓ_k only; the formula is

$$\mathbf{c}(d\theta'_k) - [\mathbb{D}, \mathbf{c}(\theta'_k)] = 2 \sum_{j=1}^n \mathbf{c}(\iota_{e_j} \theta'_k) \nabla_{e_j} - \mathbf{c}(d^* \theta'_k),$$

in terms of a local basis e_1, \dots, e_n of TX , where d^* is the codifferential). This means that only term involving θ''_k of (9.4) is relevant for calculation of the short-time limit. The term

$$F(\theta_k, \theta_{k+1}) = (-1)^{|\theta_k|} (\mathbf{c}(\theta_k) \mathbf{c}(\theta_{k+1}) - \mathbf{c}(\theta_k \wedge \theta_{k+1}))$$

has degree $\ell_k + \ell_{k+1}$ in the Getzler filtration. This shows that the term of highest order in the Getzler on the right hand side of (9.3) is the summand with $M = N$; it has order $\ell_0 + \dots + \ell_N + N$. Using the explicit formula (4.9) for Φ_T and the previous observation that only the terms involving θ''_k contribute for $k \geq 1$, we therefore obtain that modulo higher order in T , we have

$$\text{Str}(\mathbf{c}(\theta'_0) \Phi_T^{\mathcal{M}_{\mathbb{T}}^X}(\theta_1, \dots, \theta_N)) \sim (-T)^N \text{Str}(\mathbf{c}(\theta'_0) \{-\mathbf{c}(\theta''_1), \dots, -\mathbf{c}(\theta''_N)\}_{Q_T}).$$

Getzler's technique now allows to compute this term explicitly, up to higher order in T ; the result is

$$\text{Str}\left(\mathbf{c}(\theta'_0)\{\mathbf{c}(\theta''_1), \dots, \mathbf{c}(\theta''_N)\}_{Q_T}\right) \sim \frac{T^{-m_N/2-N}}{(2\pi i)^{n/2}N!} \int_X \widehat{A}(X) \wedge \theta'_0 \wedge \theta''_1 \wedge \dots \wedge \theta''_N,$$

where the additional $N!$ in the denominator comes from evaluating the integral over Δ_N contained in the bracket $\{\dots\}_{Q_T}$ (c.f. Thm. 4.2 in [7] and [16]). Substituting this into (9.3) shows that indeed, the small- T -limit is given by $\mu_0(\theta_0, \dots, \theta_N)$. \square

With this result at hand, we now use our theory to construct the desired path integral map I , which will be an integration functional for differential forms on the loop space LX . In other words, it will be a linear functional, defined on a suitable space of *integrable differential forms* on LX .

Our theory connects to the loop space through Chen's iterated integral map [18, 19], or rather its extended version, constructed by Getzler, Jones and Petrack [34], which is a map

$$\rho : \mathcal{C}(\Omega_{\mathbb{T}}(X)) \longrightarrow \Omega(LX) = \bigoplus_{k=0}^{\infty} \Omega^k(LX). \quad (9.5)$$

In order to provide a definition that is both concise and explicit, given $\theta \in \Omega^k(X)$ and $\tau \in S^1$ we denote by $\theta(\tau) \in \Omega^k(LX)$ the differential form defined at $\gamma \in LX$ by the formula

$$\theta(\tau)_\gamma[v_1, \dots, v_k] = \theta_{\gamma(\tau)}[v_1(\tau), \dots, v_k(\tau)],$$

for $v_1, \dots, v_k \in T_\gamma LX = C^\infty(S^1, \gamma^*TX)$. The (extended) iterated integral map is then given by the formula¹⁰

$$\rho(\theta_0, \dots, \theta_N) = \int_{\Delta_N} \theta'_0(0) \wedge (\iota_K \theta'_1(\tau_1) + \theta''(\tau_1)) \wedge \dots \wedge (\iota_K \theta'_N(\tau_N) + \theta''(\tau_N)) d\tau,$$

where $K(\gamma) = \dot{\gamma}$ denotes the canonical velocity vector field on LX .

Lemma 9.3 (Properties of ρ). *The extended iterated integral map is degree-preserving and satisfies*

$$\rho(\underline{d} + \underline{b}) = d\rho, \quad \rho \underline{B} - \mathbf{A} \rho \underline{L} = \mathbf{A} \iota_K \rho, \quad (9.6)$$

where \mathbf{A} is the operator on $\Omega(LX)$ that averages over the S^1 -action. Moreover, if X is connected, we have $\ker(\rho) \subseteq \ker(\text{Ch}(\mathcal{M}^X))$.

Proof. The identities (9.6) follow from tedious, but rather straight-forward calculations; c.f. the proof of Prop. 4.1 in [17]. Now if X is connected, it turns out that the kernel of ρ is precisely the subcomplex $\mathbf{D}^{\mathbb{T}}(\Omega)$ defined in (3.12), so the Lemma follows from the fact that $\text{Ch}(\mathcal{M}_{\mathbb{T}}^X)$ is Chen normalized. The inclusion $\mathbf{D}^{\mathbb{T}}(\Omega) \subseteq \ker(\rho)$ is a simple calculation; the converse inclusion is a generalization of Lemma 4.1 of [18], which follows from observations similar to those made in [34, §2]. \square

¹⁰The sign difference compared to [34, 17] is needed in order to compensate for different sign conventions of the differentials, to ensure that (9.6) still remains true.

Assume from now on that X is connected; then these results on ρ allow to define the desired integration functional for differential forms on the loop space. It is defined on the subclass of differential forms which can be represented by iterated integrals, in other words, on the subspace $\text{im}(\rho) \subset \Omega(\text{LX})$. The definition is

$$I[\xi] := i^{n/2} \langle \text{Ch}(\mathcal{M}_{\mathbb{T}}^X), c \rangle, \quad \text{if } \xi = \rho(c) \text{ for } c \in \mathbf{C}(\Omega(X)_{\mathbb{T}}), \quad (9.7)$$

where the factor $i^{n/2}$ ensures that I is real-valued on real forms ξ . Notice that if ξ is represented as an iterated integral in two ways, say by $c, c' \in \mathbf{C}(\Omega(X)_{\mathbb{T}})$, then one has $c - c' \in \ker(\rho)$ and Lemma 9.3 implies that $\langle \text{Ch}(\mathcal{M}_{\mathbb{T}}^X), c - c' \rangle = 0$, as $\text{Ch}(\mathcal{M}_{\mathbb{T}}^X)$ is Chen normalized. In other words, $I[\xi]$ is independent of the choice of c . From (9.6), it follows the integral map satisfies

$$I[(d + \iota_K)\xi] = 0, \quad (9.8)$$

which can be seen as a version of *Stokes theorem*. Indeed, the graded cyclicity of the supertrace implies that $I[\xi] = I[\mathbf{A}\xi]$ for all iterated integrals ξ (not that \mathbf{A} preserves $\text{im}(\rho)$). From (9.6) then follows that the *equivariant iterated integral map* $\mathbf{A}\rho$ sends $\underline{d}_{\mathbb{T}} + \underline{b} + \underline{B}$ to $d + \iota_K$, so that (9.8) follows from the closedness of the Chern character, Thm. 5.3.

Remark 9.4. In several places, for example [17, 34], the equivariant differential $d + P$ is used instead, where $P = \mathbf{A}\iota_K$. This difference is immaterial in our considerations, as I is cyclically invariant, $I[\mathbf{A}\xi] = I[\xi]$. Therefore, the Stokes property (9.8) holds also with P instead of ι_K , and Thm. 9.5 also holds for all iterated integrals ξ with $(d + P)\xi = 0$.

Particularly interesting integrands for the path integral map are the *Bismut-Chern characters* $\text{BCh}(E, \nabla)$ defined in [10], which are equivariantly closed differential forms on LX associated to a vector bundle E with connection ∇ over X . However, they are not contained in the algebra $\Omega(\text{LX})$, as they are an infinite sum of their homogenous components. To extend the iterated integral map to this context, we use Prop. 4.1 in [17] which implies that the iterated integral map (9.5) extends by continuity to a map

$$\rho_{\epsilon} : \mathbf{C}^{\epsilon}(\Omega(X)_{\mathbb{T}}) \longrightarrow \widehat{\Omega}(\text{LX}) := \prod_{k=0}^{\infty} \Omega^k(\text{LX}),$$

where $\Omega^k(\text{LX})$ is equipped with its canonical locally convex topology and $\widehat{\Omega}(\text{LX})$ with the induced product topology. Here, the locally convex topology on $\Omega^k(\text{LX})$ is given by the family of seminorms $\nu_f(\xi) := \nu(f^*\xi)$, where f is a smooth map from a finite dimensional manifold Y to LX and ν is a continuous seminorm on $\Omega^k(Y)$ with respect to its natural Fréchet topology. This is much in the spirit of Chen's theory of diffeologies [18].

Now using ρ_{ϵ} instead of ρ allows to extend the definition (9.7) of I to the differential forms ξ that are *entire* iterated integrals, i.e. that lie in the image of ρ_{ϵ} , so that with

$$\Omega_{\text{int}}(\text{LX}) := \text{im}(\rho_{\epsilon}) \subset \widehat{\Omega}(\text{LX})$$

the space of *integrable differential forms* on LX , the linear functional I given by (9.7) extends to a linear functional on $\Omega_{\text{int}}(\text{LX})$.

Theorem 9.5 (Fundamental properties of I). *Let X be a compact even-dimensional connected spin manifold. Then the linear functional*

$$I : \Omega_{\text{int}}(\mathbf{L}X) \longrightarrow \mathbb{C}$$

defined above is even and supersymmetric. Moreover, for all differential forms $\xi \in \Omega_{\text{int}}(\mathbf{L}X)$ with $(d + \iota_K)\xi = 0$, one has the localization formula

$$I[\xi] = (2\pi)^{-n/2} \int_X \widehat{A}(X) \wedge \xi|_X. \quad (9.9)$$

We remark that by the observations of Atiyah [5], this is precisely the formula one gets when formally applying the finite-dimensional localization formula, c.f. e.g. [25, 8] or [7, Thm. 7.13].

Proof. That I is even follows from the corresponding property of the Chern character; the supersymmetry was already noted in (9.8) and, via Stokes theorem on X , is in fact also a consequence of (9.9). It therefore remains to prove the localization formula, which follows essentially from Thm. 9.1. Indeed, let $\xi = \rho_\epsilon(c)$ for $c \in \mathbf{C}^\epsilon(\Omega(X)_\mathbb{T})$, so that

$$I[\xi] = \langle \text{Ch}(\mathcal{M}_\mathbb{T}^X), c \rangle = \langle \text{Ch}(\mathcal{M}_\mathbb{T}^X) - \mu_0, c \rangle + \langle \mu_0, c \rangle.$$

Now by Thm. 9.1, $\text{Ch}(\mathcal{M}_\mathbb{T}^X)$ and μ_0 are cohomologous, in other words, there exists a Chen normalized cochain $\mu' \in \mathbf{N}_{\mathbb{T}, \alpha}^-(\Omega)$ such that $\text{Ch}(\mathcal{M}_\mathbb{T}^X) - \mu_0 = (\underline{d} + \underline{b} + \underline{B})\mu'$. Therefore

$$\langle \text{Ch}(\mathcal{M}_\mathbb{T}^X) - \mu_0, c \rangle = -\langle \mu', (\underline{d}_\mathbb{T} + \underline{b} + \underline{B})c \rangle. \quad (9.10)$$

However, since ξ is closed, Lemma 9.3 implies that $(\underline{d} + \underline{b} + \underline{B})c \in \mathbf{D}^\mathbb{T}(\Omega)$, which shows that the right hand side of (9.10) is zero, as μ' is Chen normalized. We therefore have $I[\xi] = \langle \mu_0, c \rangle$, and the result follows after observing that

$$\rho(\theta_0, \dots, \theta_N)|_X = \frac{1}{N!} \theta'_0 \wedge \theta''_1 \wedge \dots \wedge \theta''_N,$$

where the factor $1/N!$ comes from the integral over Δ_N in the definition of ρ . □

To see how this connects to the twisted Atiyah-Singer index theorem, let (E, ∇) be a vector bundle with connection on X and let $\text{BCh}(E, \nabla)$ be the corresponding Bismut-Chern character, as defined in [10]. Then by the results of [34], one has $\text{BCh}(E, \nabla) = \rho_\epsilon(\text{Ch}(p))$ for a suitable idempotent p , hence a corollary of Thm. 8.1 is the formula

$$I[\text{BCh}(E, \nabla)] = i^{n/2} \langle \text{Ch}(\mathcal{M}_\mathbb{T}^X), \text{Ch}(p) \rangle = i^{n/2} \text{ind}(\mathbf{D}^E), \quad (9.11)$$

where \mathbf{D}^E is the Dirac operator twisted by (E, ∇) . On the other hand, the Bismut-Chern characters are equivariantly closed, hence by Thm. 9.5, we have

$$I[\text{BCh}(E, \nabla)] = (2\pi)^{-n/2} \int_X \widehat{A}(X) \wedge \text{BCh}(E, \nabla)|_X. \quad (9.12)$$

The Bismut-Chern characters now have the property that $\text{BCh}(E, \nabla) = \text{Ch}(E, \nabla)$, the usual Chern character on X of (E, ∇) , defined using Chern-Weil theory. Hence combining (9.12) and (9.11), we get the Atiyah-Singer index formula

$$\text{ind}(D^E) = (2\pi i)^{-n/2} \int_X \widehat{A}(X) \wedge \text{Ch}(E, \nabla)$$

for the twisted Dirac operator.

This puts the formal arguments of [5, 10] on a solid mathematical ground and (together with the results of [36, 37]) justifies to call I the *supersymmetric path integral* associated to X .

References

- [1] L. Alvarez-Gaumé. Supersymmetry and the Atiyah-Singer index theorem. *Comm. Math. Phys.*, 90(2):161–173, 1983.
- [2] L. Alvarez-Gaumé. Supersymmetry and index theory. In *Supersymmetry (Bonn, 1984)*, volume 125 of *NATO Adv. Sci. Inst. Ser. B Phys.*, pages 1–44. Plenum, New York, 1985.
- [3] L. Andersson and B. K. Driver. Finite-dimensional approximations to Wiener measure and path integral formulas on manifolds. *J. Funct. Anal.*, 165(2):430–498, 1999.
- [4] M. F. Atiyah. Global theory of elliptic operators. In *Proc. Internat. Conf. on Functional Analysis and Related Topics (Tokyo, 1969)*, pages 21–30. Univ. of Tokyo Press, Tokyo, 1970.
- [5] M. F. Atiyah. Circular symmetry and stationary-phase approximation. *Astérisque*, 131:43–59, 1985. Colloquium in honor of Laurent Schwartz, Vol. 1 (Palaiseau, 1983).
- [6] F. Baudoin. Brownian Chen series and Atiyah-Singer theorem. *J. Funct. Anal.*, 254(2):301–317, 2008.
- [7] N. Berline, E. Getzler, and M. Vergne. *Heat kernels and Dirac operators*. Grundlehren Text Editions. Springer-Verlag, Berlin, 2004. Corrected reprint of the 1992 original.
- [8] N. Berline and M. Vergne. Classes caractéristiques équivariantes. Formule de localisation en cohomologie équivariante. *C. R. Acad. Sci. Paris Sér. I Math.*, 295(9):539–541, 1982.
- [9] J.-M. Bismut. The Atiyah-Singer theorems: a probabilistic approach. I. The index theorem. *J. Funct. Anal.*, 57(1):56–99, 1984.
- [10] J.-M. Bismut. Index theorem and equivariant cohomology on the loop space. *Comm. Math. Phys.*, 98(2):213–237, 1985.

- [11] J.-M. Bismut. Localization formulas, superconnections, and the index theorem for families. *Comm. Math. Phys.*, 103(1):127–166, 1986.
- [12] J.-M. Bismut. The hypoelliptic Laplacian on the cotangent bundle. *J. Amer. Math. Soc.*, 18(2):379–476, 2005.
- [13] J.-M. Bismut. The hypoelliptic Dirac operator. In *Geometry and dynamics of groups and spaces*, volume 265 of *Progr. Math.*, pages 113–246. Birkhäuser, Basel, 2008.
- [14] J.-M. Bismut. *Hypoelliptic Laplacian and orbital integrals*, volume 177 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2011.
- [15] B. Blackadar. *K-theory for operator algebras*, volume 5 of *Mathematical Sciences Research Institute Publications*. Cambridge University Press, Cambridge, second edition, 1998.
- [16] J. Block and J. Fox. Asymptotic pseudodifferential operators and index theory. In *Geometric and topological invariants of elliptic operators (Brunswick, ME, 1988)*, volume 105 of *Contemp. Math.*, pages 1–32. Amer. Math. Soc., Providence, RI, 1990.
- [17] S. Cacciatori and B. Güneysu. Odd characteristic classes in entire cyclic homology and equivariant loop space homology. *ArXiv e-prints*, May 2018. To appear in *J. Noncommut. Geom.*
- [18] K. T. Chen. Iterated integrals of differential forms and loop space homology. *Ann. of Math. (2)*, 97:217–246, 1973.
- [19] K. T. Chen. Reduced bar constructions on de Rham complexes. In *Algebra, topology, and category theory (a collection of papers in honor of Samuel Eilenberg)*, pages 19–32. Academic Press, New York, 1976.
- [20] K. T. Chen. Reduced bar constructions on de Rham complexes. In *Algebra, topology, and category theory (a collection of papers in honor of Samuel Eilenberg)*, pages 19–32. Academic Press, New York, 1976.
- [21] A. Connes. Entire cyclic cohomology of Banach algebras and characters of θ -summable Fredholm modules. *K-Theory*, 1(6):519–548, 1988.
- [22] A. Connes. Compact metric spaces, Fredholm modules, and hyperfiniteness. *Ergodic Theory Dynam. Systems*, 9(2):207–220, 1989.
- [23] A. Connes and H. Moscovici. Cyclic cohomology, the Novikov conjecture and hyperbolic groups. *Topology*, 29(3):345–388, 1990.
- [24] A. Connes and H. Moscovici. The local index formula in noncommutative geometry. *Geom. Funct. Anal.*, 5(2):174–243, 1995.

- [25] J. J. Duistermaat and G. J. Heckman. On the variation in the cohomology of the symplectic form of the reduced phase space. *Invent. Math.*, 69(2):259–268, 1982.
- [26] D. S. Fine and S. Sawin. Path integrals, supersymmetric quantum mechanics, and the Atiyah-Singer index theorem for twisted Dirac. *J. Math. Phys.*, 58(1):012102, 30, 2017.
- [27] D. S. Fine and S. F. Sawin. A rigorous path integral for supersymmetric quantum mechanics and the heat kernel. *Comm. Math. Phys.*, 284(1):79–91, 2008.
- [28] D. Friedan and P. Windey. Supersymmetric derivation of the Atiyah-Singer index and the chiral anomaly. *Nuclear Phys. B*, 235(3):395–416, 1984.
- [29] E. Getzler. A short proof of the local Atiyah-Singer index theorem. *Topology*, 25(1):111–117, 1986.
- [30] E. Getzler. Path integral of the Dirac operator. *Unpublished*, 1988.
- [31] E. Getzler. The Thom class of Mathai and Quillen and probability theory. In *Stochastic analysis and applications (Lisbon, 1989)*, volume 26 of *Progr. Probab.*, pages 111–122. Birkhäuser Boston, Boston, MA, 1991.
- [32] E. Getzler. The odd Chern character in cyclic homology and spectral flow. *Topology*, 32(3):489–507, 1993.
- [33] E. Getzler and J. D. S. Jones. A_∞ -algebras and the cyclic bar complex. *Illinois J. Math.*, 34(2):256–283, 1990.
- [34] E. Getzler, J. D. S. Jones, and S. Petrack. Differential forms on loop spaces and the cyclic bar complex. *Topology*, 30(3):339–371, 1991.
- [35] E. Getzler and A. Szenes. On the Chern character of a theta-summable Fredholm module. *J. Funct. Anal.*, 84(2):343–357, 1989.
- [36] F. Hanisch and M. Ludewig. Supersymmetric path integrals I: Differential forms on the loop space. *ArXiv e-prints*, Sept. 2017.
- [37] F. Hanisch and M. Ludewig. Supersymmetric path integrals II: The fermionic integral and pfaffian line bundles. *ArXiv e-prints*, Sept. 2017.
- [38] N. Higson. The local index formula in noncommutative geometry. In *Contemporary developments in algebraic K-theory*, ICTP Lect. Notes, XV, pages 443–536. Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004.
- [39] A. Jaffe, A. Lesniewski, and K. Osterwalder. Quantum K -theory. I. The Chern character. *Comm. Math. Phys.*, 118(1):1–14, 1988.

- [40] J. D. S. Jones and R. Léandre. L^p -Chen forms on loop spaces. In *Stochastic analysis (Durham, 1990)*, volume 167 of *London Math. Soc. Lecture Note Ser.*, pages 103–162. Cambridge Univ. Press, Cambridge, 1991.
- [41] R. Léandre. Cohomologie de Bismut-Nualart-Pardoux et cohomologie de Hochschild entière. In *Séminaire de Probabilités, XXX*, volume 1626 of *Lecture Notes in Math.*, pages 68–99. Springer, Berlin, 1996.
- [42] J.-L. Loday. *Cyclic homology*, volume 301 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1992. Appendix E by María O. Ronco.
- [43] J. Lott. Supersymmetric path integrals. *Comm. Math. Phys.*, 108(4):605–629, 1987.
- [44] R. Meyer. *Local and analytic cyclic homology*, volume 3 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2007.
- [45] M. S. Narasimhan and S. Ramanan. Existence of universal connections. *Amer. J. Math.*, 83:563–572, 1961.
- [46] D. Perrot. A bivariant Chern character for families of spectral triples. *Comm. Math. Phys.*, 231(1):45–95, 2002.
- [47] D. Quillen. Algebra cochains and cyclic cohomology. *Inst. Hautes Études Sci. Publ. Math.*, 68:139–174 (1989), 1988.
- [48] S. Stolz and P. Teichner. The spinor bundle on the loop space. https://people.mpim-bonn.mpg.de/teichner/Math/Surveys_files/MPI.pdf, 2005.
- [49] S. Stolz and P. Teichner. Supersymmetric field theories and generalized cohomology. In *Mathematical foundations of quantum field theory and perturbative string theory*, volume 83 of *Proc. Sympos. Pure Math.*, pages 279–340. Amer. Math. Soc., Providence, RI, 2011.
- [50] K. Waldorf. Spin structures on loop spaces that characterize string manifolds. *Algebr. Geom. Topol.*, 16(2):675–709, 2016.
- [51] S. O. Wilson. A loop group extension of the odd Chern character. *J. Geom. Phys.*, 102:32–43, 2016.
- [52] E. Witten. Constraints on supersymmetry breaking. *Nuclear Phys. B*, 202(2):253–316, 1982.
- [53] E. Witten. Supersymmetry and Morse theory. *J. Differential Geom.*, 17(4):661–692 (1983), 1982.