

Kac regular sets and Sobolev spaces in geometry, probability and quantum physics

Francesco Bei and Batu Güneysu

Francesco Bei, Dipartimento di matematica, universit a degli studi di Padova, Italy
E-mail address: bei@math.unipd.it

Batu Güneysu, Mathematisches Institut, Universit at Bonn, Germany
E-mail address: gueneysu@math.uni-bonn.de

Abstract. Let $\Omega \subset M$ be an open subset of a Riemannian manifold M and let $V : M \rightarrow \mathbb{R}$ be a Kato decomposable potential. With $W_0^{1,2}(M; V)$ the natural form domain of the Schrödinger operator $-\Delta + V$ in $L^2(M)$, in this paper we study systematically the following question: Under which assumption on Ω is the statement

for all $f \in W_0^{1,2}(M; V)$ with $f = 0$ a.e. in $M \setminus \Omega$ one has $f|_\Omega \in W_0^{1,2}(\Omega; V)$

true for every such V ? We prove that without any further assumptions on V , the above property is satisfied, if Ω is Kac regular, a probabilistic property which means that the first exit time of Brownian motion on M from Ω is equal to its first penetration time to $M \setminus \Omega$. In fact, we treat more general covariant Schrödinger operators acting on sections in metric vector bundles, allowing new results concerning the harmonicity of Dirac spinors on singular subsets. Finally, we prove that Lipschitz regular Ω 's are Kac regular.

1. Introduction

Consider the following three properties that an open subset Ω of a noncompact Riemannian manifold M may or may not have:

(A) Let V be a Kato decomposable potential on M and let $H_M(V)$ denote the natural quadratic form realization of $-\Delta + V$ in $L^2(M)$ and let $H_\Omega(V)$ denote the self-adjoint realization of $-\Delta + V$ in $L^2(\Omega)$ subject to Dirichlet boundary conditions (also the Friedrichs realization). *For every such V one has*

$$(1) \quad \exp(-t(H_M(V) + \infty \cdot 1_{M \setminus \Omega})) = \exp(-tH_\Omega(V))P_\Omega \quad \text{for all } t > 0,$$

where $\exp(-t(H_M(V) + \infty \cdot 1_{M \setminus \Omega}))$ is understood as the strong limit of $\exp(-t(H_M(V) + n \cdot 1_{M \setminus \Omega}))$ as $n \rightarrow \infty$, and where $P_\Omega : L^2(M) \rightarrow L^2(\Omega)$ denotes the natural projection.

(B) Assume M is a geodesically complete Riemannian spin manifold. *For every spin bundle $\mathcal{S} \rightarrow M$ with corresponding Dirac operator D acting on sections of $\mathcal{S} \rightarrow M$, and every*

spinor $\Psi \in \Gamma_{L^2}(M, \mathcal{S})$ with $D\Psi \in \Gamma_{L^2}(M, \mathcal{S})$ and $\Psi = 0$ almost everywhere in $M \setminus \Omega$, one has the implication

$$D^2\Psi = 0 \text{ in } \Omega \Rightarrow D\Psi = 0 \text{ in } \Omega .$$

(C) The first exit time of Brownian motion from Ω equals the first penetration time of Brownian motion to $M \setminus \Omega$. This property in particular implies¹ that with \mathbb{P}^x the Riemannian Brownian motion measure with initial point x and \mathbb{X} the coordinate process on the path space of M one has

$$\begin{aligned} & \{\mathbb{X}_s \in \Omega \text{ for a.e. } s \in [0, t] \text{ and } \mathbb{X}_s \in M \text{ for all } s \in [0, t]\} \\ & \stackrel{=_{\mathbb{P}^x}}{\text{}} \{\mathbb{X}_s \in \Omega \text{ for all } s \in [0, t]\} \quad \text{for all } x \in \Omega, t > 0. \end{aligned}$$

To the best of our knowledge, property (C) has been introduced by D. Stroock in the Euclidean space \mathbb{R}^m , and is usually referred to as the *Kac regularity* of Ω .

While (A), (B) and (C) seem unrelated at first glance, the main results of this paper show that the above three problems are much more correlated than one might expect: Indeed we prove that given an arbitrary open subset $\Omega \subset M$,

(i) (A) is equivalent to

(A') for all $f \in W_0^{1,2}(M; V)$ with $f = 0$ a.e. in $M \setminus \Omega$ one has $f|_{\Omega} \in W_0^{1,2}(\Omega; V)$

for every V as in (A); here $W_0^{1,2}(M; V)$ and $W_0^{1,2}(\Omega; V)$ denote, respectively, the form domain of $H_M(V)$ and $H_{\Omega}(V)$,

(ii) (C) is equivalent to (A),

(iii) (C) implies (B),

(iv) If Ω is Lipschitz regular (cf. Definition 2.13), then one has (C).

While (i), (ii) and a variant of (iv) have been obtained in the Euclidean space \mathbb{R}^m [13] for $V = 0$ (cf. Remark 2.16), our analysis is the first systematic treatment of these questions on manifolds, allowing in addition potentials and in fact covariant Schrödinger operators. As we only require a very weak regularity on the potential V , we can also treat Schrödinger operators that appear naturally in quantum mechanics, having potentials with Coulomb type singularities. In fact, we are going to treat these problems within the much more general class of covariant Schrödinger operators that act on sections of metric vector bundles over M , allowing to treat magnetic fields or squares of geometric Dirac operators simulatinously, making contact with (B). Our proofs rely on various (partially new) covariant Feynman-Kac formulas.

2. Main result

In the sequel, we work in the smooth category, that is, all differential, topological and geometric data (like manifolds, bundles, metrics and covariant derivatives) are understood to be smooth. In addition, any manifold is understood to be without boundary, unless otherwise stated. Without loss of generality, we will consider only complex function spaces.

¹We will not assume M to be stochastically complete

Furthermore, for every manifold M and every complex vector bundle $E \rightarrow M$ its fibers will be denoted with E_x , $x \in M$, and the corresponding space of sections of $E \rightarrow M$ having a certain regularity \mathcal{C} like $\mathcal{C} = C_c^\infty$ etc. will be denoted with $\Gamma_{\mathcal{C}}(M, E)$. The smooth linear partial differential operator

$$d_\bullet : \Gamma_{C^\infty}(M, \wedge^\bullet T_{\mathbb{C}}^* M) \longrightarrow \Gamma_{C^\infty}(M, \wedge^{\bullet+1} T_{\mathbb{C}}^* M)$$

from $\wedge^\bullet T_{\mathbb{C}}^* M \rightarrow M$ to itself denotes the exterior derivative acting on complexified differential forms.

Let M be an arbitrary connected Riemannian m -manifold. As such, M is equipped with its Riemannian volume measure μ . Given a complex metric vector bundle $E \rightarrow M$, the scalar product on the complex Hilbert space $\Gamma_{L^2}(M, E)$ of Borel equivalence classes of square integrable sections of $E \rightarrow M$ is simply denoted with

$$\langle f_1, f_2 \rangle = \int (f_1, f_2) d\mu,$$

with

$$\|f\|^2 = \int |f|^2 d\mu$$

the induced norm. Given two such bundles $E_1 \rightarrow M$, $E_2 \rightarrow M$, the formal adjoint of a smooth linear partial differential operator D mapping sections of $E_1 \rightarrow M$ to sections of $E_2 \rightarrow M$ with respect to $\langle \cdot, \cdot \rangle$ is simply denoted with D^\dagger . The reader may find the basics of linear partial differential operators on vector bundles over Riemannian manifolds in [11]. The following conventions will be very convenient in the sequel:

Notation 2.1. If $\Omega \subset M$ is an open subset and $E \rightarrow M$ is a complex vector bundle, we define

$$\Gamma_{C_c^\infty}(\Omega, E) := \{f \in \Gamma_{C_c^\infty}(M, E) : f \text{ is compactly supported in } \Omega\} \subset \Gamma_{C_c^\infty}(M, E),$$

and if $E \rightarrow M$ is equipped with a metric, then

$$\Gamma_{L^2}(\Omega, E) := \{f \in \Gamma_{L^2}(M, E) : f = 0 \text{ } \mu\text{-a.e. in } M \setminus \Omega\} \subset \Gamma_{L^2}(M, E).$$

Then $\Gamma_{L^2}(\Omega, E)$ is a closed subspace of $\Gamma_{L^2}(M, E)$, thus a Hilbert space in itself. We denote with

$$P_\Omega : \Gamma_{L^2}(M, E) \longrightarrow \Gamma_{L^2}(\Omega, E)$$

the orthogonal projection onto $\Gamma_{L^2}(\Omega, E)$. Note that P_Ω is nothing but the restriction map $f \mapsto f|_\Omega$.

Let $(t, y) \mapsto p_\Omega(t, x, y)$ denote the pointwise minimal solution of the heat equation on the open subset $\Omega \subset M$ with $\lim_{t \rightarrow 0^+} p_\Omega(t, x, \cdot) = \delta_x$. We recall that the Kato class $\mathcal{K}(M)$ of M is defined by all Borel functions w on M such that

$$(2) \quad \lim_{t \rightarrow 0^+} \sup_{x \in M} \int p_M(t, x, y) |w(y)| d\mu(y) = 0.$$

Based on the results from [11] (cf. Chapter VII), we propose:

Definition 2.2. By a *Kato-Schrödinger bundle over M* , we will understand a datum

$$(E, \nabla, V) \longrightarrow M$$

with

- $E \rightarrow M$ a complex metric vector bundle
- ∇ a metric covariant derivative on $E \rightarrow M$,
- $V : M \rightarrow \text{End}(E)$ is a Kato decomposable potential, that is, there exist pointwise self-adjoint Borel sections V_{\pm} of $\text{End}(E) \rightarrow M$ with $V_{\pm}(x) \geq 0$ for all $x \in M$ and

$$(3) \quad V = V_+ - V_-, \quad |V_+| \in L_{\text{loc}}^1(M), \quad |V_-| \in \mathcal{K}(M),$$

where $|\cdot|$ denotes the fiberwise operator norm.

It follows in the above situation that $|V_-| \in L_{\text{loc}}^1(M)$ (cf. Lemma VI.3 in [11]) and so $|V| \in L_{\text{loc}}^1(M)$. In view of [7]

$$\int p_M(t, x, y) d\mu(y) \leq 1 \quad \text{for all } (t, x) \in (0, \infty) \times M,$$

it follows that every bounded Borel function w on M satisfies (2), and the reader may find various (weighted) L^q -assumptions on w that imply (2) in [19, 20, 8] (see also Chapter VI from [11]). In particular, every pointwise self-adjoint L_{loc}^1 -section V of $\text{End}(E) \rightarrow M$ which is bounded from below, in the sense that for some constant $C \in \mathbb{R}$ one has $V(x) \geq C$ for all $x \in M$, satisfies (3).

Notation 2.3. In the situation of Definition 2.2, let $\Omega \subset M$ be an open subset. Then $\Gamma_{W_0^{1,2}}(\Omega, E; \nabla, V)$ is defined to be the space of all $f \in \Gamma_{L^2}(\Omega, E; \nabla, V)$ which admit a sequence $(f_n) \subset \Gamma_{C^\infty}(\Omega, E; \nabla, V)$ such that

$$\|f_n - f\| \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

$$\int (\nabla^\dagger \nabla (f_n - f_m), (f_n - f_m)) d\mu + \int (V(f_n - f_m), (f_n - f_m)) d\mu \rightarrow 0 \quad \text{as } n, m \rightarrow \infty.$$

By Corollary XIII.3 in [11] the symmetric densely defined sesquilinear form

$$\Gamma_{C_c^\infty}(\Omega, E) \times \Gamma_{C_c^\infty}(\Omega, E) \ni (f_1, f_2) \longmapsto \int (\nabla^\dagger \nabla f_1, f_2) d\mu + \int (V f_1, f_2) d\mu \in \mathbb{C}$$

is closable and semibounded from below in $\Gamma_{L^2}(\Omega, E)$. Its closure $\langle \cdot, \cdot \rangle_{\nabla, V, *}$ has the domain of definition $\Gamma_{W_0^{1,2}}(\Omega, E; \nabla, V)$ and is explicitly given as follows: For $f, \tilde{f} \in \Gamma_{W_0^{1,2}}(\Omega, E; \nabla, V)$ with defining sequences $(f_n), (\tilde{f}_n) \subset \Gamma_{C_c^\infty}(\Omega, E; \nabla, V)$, one has

$$\langle f, \tilde{f} \rangle_{\nabla, V, *} = \lim_{n \rightarrow \infty} \left(\int (\nabla^\dagger \nabla f_n, \tilde{f}_n) d\mu + \int (V f_n, \tilde{f}_n) d\mu \right),$$

and this number does not depend on the defining sequences. The semibounded from below and self-adjoint operator in $\Gamma_{L^2}(\Omega, E)$ induced by $\langle \cdot, \cdot \rangle_{\nabla, V, *}$ is denoted with $H_\Omega(\nabla, V)$. In addition, $\Gamma_{W_0^{1,2}}(\Omega, E; \nabla, V)$ becomes a Hilbert space with respect to the scalar product

$$\langle f_1, f_2 \rangle_{\nabla, V} := (1 + C) \langle f_1, f_2 \rangle + \langle f_1, f_2 \rangle_{\nabla, V, *},$$

where $C \geq 0$ is any constant satisfying

$$(4) \quad \int (\nabla^\dagger \nabla f, f) d\mu + \int (Vf, f) d\mu \geq -C \|f\|^2 \quad \text{for all } f \in \Gamma_{C^\infty}(M, E),$$

noting that any two such constants produce equivalent scalar products.

Note also that $H_\Omega(\nabla, V)$ is nothing but the Friedrichs realization of $\nabla^\dagger \nabla + V$ in $\Gamma_{L^2}(\Omega, E)$. Traditionally, if $\Omega \neq M$, one says that $H_\Omega(\nabla, V)$ is the self-adjoint realization of $\nabla^\dagger \nabla + V$ in $\Gamma_{L^2}(\Omega, E)$ subject to Dirichlet boundary conditions.

To simplify the notation in some special cases, we add:

Remark 2.4. If $E \rightarrow M$ is the trivial complex line bundle and ∇ the covariant derivative which is induced by the exterior differential, then we simply omit ∇ and E everywhere in the notation. Note that in this case $\nabla^\dagger \nabla =: -\Delta$ is by definition the usual Laplace-Beltrami operator on functions. In case $V = 0$ we will in addition omit V everywhere in the notation. These conventions lead to natural notations such as $W_0^{1,2}(\Omega; V)$, $H_\Omega(V)$, $W_0^{1,2}(\Omega)$, $L^2(\Omega)$, H_Ω . In particular, the self-adjoint operator H_Ω in $L^2(\Omega)$ is $-\Delta$ with Dirichlet boundary conditions and the integral kernel $\exp(-tH_\Omega)(x, y)$ of $\exp(-tH_\Omega)$ is precisely $p_\Omega(t, x, y)$ [7].

Here is an example from quantum mechanics:

Example 2.5. If M is the Euclidean space \mathbb{R}^3 and $\eta = \sum_{j=1}^3 \eta_j dx^j$ is a smooth real-valued 1-form on \mathbb{R}^3 (considered as a magnetic potential), $Z \in \mathbb{N}$, and $V : \mathbb{R}^3 \rightarrow \mathbb{R}$ is the Coulomb potential $V(x) := -Z/|x - x_0|$ with mass at $x_0 \in \mathbb{R}^3$, then with

$$\nabla^\eta : C^\infty(\mathbb{R}^3) \longrightarrow \Gamma_{C^\infty}(\mathbb{R}^3, T_{\mathbb{C}}^*\mathbb{R}^3), \quad \nabla^\eta f := \sum_{j=1}^3 (\partial_j f + \sqrt{-1} f \eta_j) dx^j,$$

the datum

$$(\mathbb{R}^3 \times \mathbb{C}, \nabla^\eta, V) \longrightarrow \mathbb{R}^3$$

is a Kato-Schrödinger bundle (cf. Proposition VI.14 in [11]). In this case, identifying sections in $\mathbb{R}^3 \times \mathbb{C} \rightarrow \mathbb{R}^3$ with functions on M , the semibounded self-adjoint operator $H_{\mathbb{R}^3}(\nabla^\eta, V)$ in $L^2(\mathbb{R}^3)$ is the Hamilton operator of an atom having one electron and a nucleus having Z protons, in the magnetic field given by

$$d\eta = \sum_{i < j} (\partial_i \eta_j - \partial_j \eta_i) dx^i \wedge dx^j.$$

More generally, one could replace in this example \mathbb{R}^3 with a Riemannian 3-fold M such that $p_M(t, x, y)$ satisfies a Gaussian upper bound of the form

$$p_M(t, x, y) \leq C_1 t^{-3/2} \exp\left(-\frac{d(x, y)^2}{C_2 t}\right), \quad (t, x, y) \in (0, \infty) \times M \times M,$$

taking

$$V(x) := -Z \int_0^\infty p_M(t, x, x_0) dt$$

to be the Coulomb potential with mass at x_0 . One could even take the electron's spin into account, as is shown in [10].

Here is an example from geometry:

Example 2.6. 1. A *geometric Dirac bundle over M* is understood to be a datum

$$(E; c, \nabla) \longrightarrow M$$

such that

- $E \rightarrow M$ is a complex metric vector bundle
- c is a Clifford multiplication on $E \rightarrow M$, in the sense that c is a homomorphism of real (!) vector bundles $c : TM \rightarrow \text{End}(E)$, such that for all $X \in \Gamma_{C^\infty}(M, TM)$ one has $c(X) = -c(X)^*$ and $c(X)^*c(X) = |X|^2$
- ∇ is a Clifford connection on $(E; c) \rightarrow M$, that is, ∇ is a metric covariant derivative on $E \rightarrow M$ such that for all $X, Y \in \Gamma_{C^\infty}(M, TM)$, $\Psi \in \Gamma_{C^\infty}(M, E)$ one has

$$\nabla_X(c(Y)\Psi) = c(\nabla_X^{TM}Y)\Psi + c(Y)\nabla_X\Psi.$$

Then the associated geometric Dirac operator is the first order linear partial differential operator on $E \rightarrow M$ defined by $D(c, \nabla) := \sum_{j=1}^m c(e_j)\nabla_{e_j}$, where e_j is any local orthonormal-frame for $TM \rightarrow M$ and $m = \dim M$. The operator $D(c, \nabla)$ is formally self-adjoint with symbol c , in particular elliptic. By the abstract Lichnerowicz formula [23, 21],

$$V(c, \nabla) := D(c, \nabla)^2 - \nabla^\dagger \nabla : M \rightarrow \text{End}(E)$$

is a smooth potential. In case $V(c, \nabla)$ is Kato decomposable (which is the case, e.g., if $V(c, \nabla)$ is bounded from below), we call

$$(E; c, \nabla) \longrightarrow M$$

a *geometric Kato-Dirac bundle*, and we get the Kato-Schrödinger bundle

$$(E, \nabla, V(c, \nabla)) \longrightarrow M.$$

In this case, $\Gamma_{W_0^{1,2}}(\Omega, E; \nabla, V(c, \nabla))$ is given by all $f \in \Gamma_{L^2}(\Omega, E)$ with $D(c, \nabla)f \in \Gamma_{L^2}(\Omega, E)$ in the sense of distributions, which admit a sequence $f_n \in \Gamma_{C^\infty}(\Omega, E)$, $n \in \mathbb{N}$, with

$$\|f_n - f\| + \|D(c, \nabla)f_n - D(c, \nabla)f\| \rightarrow 0.$$

We have

$$\langle f_1, f_2 \rangle_{\nabla, V(c, \nabla), *} = \langle D(c, \nabla)f_1, D(c, \nabla)f_2 \rangle,$$

so that in particular the scalar product on $\Gamma_{W_0^{1,2}}(\Omega, E; \nabla, V(c, \nabla))$ is given by

$$\langle f_1, f_2 \rangle_{\nabla, V(c, \nabla)} = \langle f_1, f_2 \rangle + \langle D(c, \nabla)f_1, D(c, \nabla)f_2 \rangle.$$

2. For example, if M is a spin manifold, then every spin structure on M canonically induces a geometric Dirac bundle over M [21, 23], which is a geometric Kato-Dirac bundle, if the scalar curvature of M is Kato decomposable.

3. Another example of a geometric Dirac bundle over M (which does not require any topological assumptions on M) is given by taking $E = \wedge T_{\mathbb{C}}^*M$ with the metric induced

by the Riemannian metric on M and ∇ the covariant derivative which is induced by the Levi-Civita connection on M , and

$$c(\alpha)\beta := \alpha \wedge \beta - \iota_{\alpha^\sharp}\beta, \quad \alpha \in \Gamma_{C^\infty}(M, T_{\mathbb{C}}^*M), \beta \in \Gamma_{C^\infty}(M, \wedge T_{\mathbb{C}}^*M),$$

where ι_{α^\sharp} denotes the contraction with the vector field α^\sharp which is induced by α via the Riemannian metric on M . In this case, one can calculate that $D(c, \nabla) = d + d^\dagger$, so that $D(c, \nabla)^2$ is by definition the Laplace-Beltrami operator acting on differential forms. Note that the restriction of $D(c, \nabla)^2$ to 0-forms is precisely the usual Laplace-Beltrami operator $-\Delta$ on functions, and that the restriction of $D(c, \nabla)$ to 0-forms can be identified with the gradient (cf. Proposition 11.2.1 in [23] for detailed proofs of these facts).

Given a Kato-Schrödinger bundle

$$(E, \nabla, V) \longrightarrow M,$$

the inclusion

$$\Gamma_{W_0^{1,2}}(\Omega, E; \nabla, V) \subset \{f \in \Gamma_{W_0^{1,2}}(M, E; \nabla, V) : f|_{M \setminus \Omega} = 0 \text{ } \mu\text{-a.e.}\}$$

is easily seen to be satisfied without any further assumptions on the underlying data: Indeed, given $\psi \in \Gamma_{W_0^{1,2}}(\Omega, E, \nabla, V)$ we can per definitionem pick a sequence $\psi_n \in \Gamma_{C_c^\infty}(\Omega, E)$ with $\|\psi_n - \psi\|_{\nabla, V} \rightarrow 0$, so that $\|\psi_n - \psi\| \rightarrow 0$ and so $\psi = 0$ μ -a.e. in $M \setminus \Omega$.

The question we address in this paper is: Under which condition on Ω does the reverse inclusion

$$(5) \quad \Gamma_{W_0^{1,2}}(\Omega, E; \nabla, V) \supset \{f \in \Gamma_{W_0^{1,2}}(M, E; \nabla, V) : f|_{M \setminus \Omega} = 0 \text{ } \mu\text{-a.e.}\}$$

hold true?

By the above, (5) is equivalent to

$$(6) \quad \Gamma_{W_0^{1,2}}(\Omega, E; \nabla, V) = \{f \in \Gamma_{W_0^{1,2}}(M, E; \nabla, V) : f|_{M \setminus \Omega} = 0 \text{ } \mu\text{-a.e.}\}.$$

The validity of (6) turns out to be equivalent to the corresponding Dirichlet heat-flow of $H_\Omega(\nabla, V)$ on Ω being realized as the heat-flow of $H_M(\nabla, V)$ on M perturbed by the ‘potential’ $\infty \cdot 1_{M \setminus \Omega}$, for one has:

Proposition 2.7. *Let $\Omega \subset M$ be an arbitrary open subset and let $(E, \nabla, V) \rightarrow M$ be a Kato-Schrödinger bundle over M . Then one has (6), if and only if for all $t \geq 0$ one has*

$$(7) \quad \lim_{n \rightarrow \infty} \exp(-tH_M(\nabla, V + n1_{M \setminus \Omega})) = \exp(-tH_\Omega(\nabla, V))P_\Omega$$

strongly as bounded operators in $\Gamma_{L^2}(M, E)$.

The proof of Proposition 2.7 will be given in Section 4. It is a consequence of monotone convergence results for sesquilinear forms.

As one might guess, the validity of (6) or (7) should depend on the local regularity of $\partial\Omega$. In order to make precise how (6) or (7) depend on the local regularity of $\partial\Omega$, the probabilistic Definition 2.10 below will turn out to be crucial. To this end, let $W(\hat{M})$ denote

the Wiener space of continuous paths $\omega : [0, \infty) \rightarrow \hat{M}$ with its Borel-sigma-algebra \mathcal{F} , where $\hat{M} = M \cup \{\infty_M\}$ is defined to be the (essentially uniquely determined) Alexandroff-compactification of M in case M is noncompact, and $\hat{M} := M$ if M is compact.² In any case, $W(\hat{M})$ is equipped with the topology of locally uniform convergence. Let

$$\mathbb{X} : [0, \infty) \times W(\hat{M}) \longrightarrow \hat{M}, \quad \mathbb{X}_t(\omega) := \omega(t)$$

denote the coordinate process, and for every $x \in M$, the symbol \mathbb{P}^x stands for the Riemannian Brownian motion measure on \mathcal{F} with

$$\mathbb{P}^x \{\mathbb{X}_0 = x\} = 1.$$

In other words, the transition density of \mathbb{P}^x with respect to μ is given by the natural extension of $\exp(-tH_M)(x, y)$ to $(0, \infty) \times \hat{M} \times \hat{M}$. We refer the reader to Section 8 in [6] and the references therein for the definition and the existence of \mathbb{P}^x .

With \mathcal{F}_* the filtration of \mathcal{F} which is generated by \mathbb{X} , the family \mathbb{P}^x , $x \in M$, has the strong Markov property for stopping times in \mathcal{F}_* . If $\Omega \subset \hat{M}$ is an open set, then³

$$\alpha_\Omega : W(\hat{M}) \longrightarrow [0, \infty], \quad \alpha_\Omega := \inf \{t > 0 : \mathbb{X}_t \in \hat{M} \setminus \Omega\}$$

denotes the first exit time of \mathbb{X} from Ω . It is well-known that α_Ω is an \mathcal{F}_* -stopping time. In particular, if M is noncompact, then the stopping time α_M is the explosion time of \mathbb{X} , and in this case the point at infinity ∞_M of \hat{M} is \mathbb{P}^x -almost surely (a.s.) absorbing for all $x \in M$, in the sense that

$$\mathbb{P}^x \left(\{\alpha_M = \infty\} \cup \{\alpha_M < \infty \text{ and } \mathbb{X}_t = \infty_M \text{ for all } t \in [\alpha_M, \infty)\} \right) = 1 \quad \text{for all } x \in M.$$

Likewise,

$$\beta_\Omega := \inf \left\{ t > 0 : \int_0^t 1_{\hat{M} \setminus \Omega}(\mathbb{X}_s) ds > 0 \right\} : W(\hat{M}) \longrightarrow [0, \infty]$$

denotes the first penetration time of \mathbb{X} into $M \setminus \Omega$. One trivially has

$$(8) \quad \alpha_\Omega \leq \beta_\Omega,$$

and again β_Ω induces an \mathcal{F}_* -stopping time. Note that both α and β are pathwise monotonely increasing in Ω , and that one pathwise has

$$\beta_\Omega = \inf \left\{ t \geq 0 : \int_0^t 1_{\hat{M} \setminus \Omega}(\mathbb{X}_s) ds > 0 \right\} = \inf \left\{ t \geq 0 : \mathbb{X}_s \in \Omega \text{ for a.e. } s \in [0, t] \right\}.$$

If $\omega \in W(\hat{M})$ is such that $\omega(0) \in \Omega$, then one also gets

$$\beta_\Omega(\omega) \geq \alpha_\Omega(\omega) = \inf \{t \geq 0 : \omega(t) \in \hat{M} \setminus \Omega\} > 0.$$

The following regularity result will be very useful in the sequel:

²Note that every open subset of M is also open in \hat{M} . In addition, the closure of a subset Ω of M is always understood with respect to the topology of M and not the topology of \hat{M} , noting that both closures coincide if and only if the closure of Ω in M is a compact subset of M .

³In the sequel the infimum of an empty set is understood to be ∞ .

Lemma 2.8. *Let $\omega \in W(\hat{M})$ be such that in case $\alpha_M(\omega) < \infty$ one has $\omega(t) = \infty_M$ for all $t \in [\alpha_M(\omega), \infty)$, let Ω be an open subset of M , and let $\Omega_n \subset \Omega$, $n \in \mathbb{N}$, be open with $\Omega_n \subset \Omega_{n+1}$ for all $n \in \mathbb{N}$, $\omega(0) \in \Omega_1$ and $\bigcup_{n \in \mathbb{N}} \Omega_n = \Omega$.*

a) *One has $\alpha_{\Omega_n}(\omega) \nearrow \alpha_\Omega(\omega)$ as $n \rightarrow \infty$.*

b) *Assume in addition that for each $n \in \mathbb{N}$ there exists an open subset $\Upsilon_n \subset M$ such that $\Upsilon_n \cap \Omega = \Omega_n$ and $\bar{\Omega} = \bigcup_{n \in \mathbb{N}} (\bar{\Omega} \cap \Upsilon_n)$. Then one has $\beta_{\Omega_n}(\omega) \nearrow \beta_\Omega(\omega)$ as $n \rightarrow \infty$.*

While Lemma 2.8 a) is well-known, a proof of Lemma 2.8 b) will be given in Section 3.

Remark 2.9. If $\{\Upsilon_n : n \in \mathbb{N}\}$ is a family of open subsets of M such that $\Upsilon_n \subset \Upsilon_{n+1}$ for all $n \in \mathbb{N}$ and $\bigcup_{n \in \mathbb{N}} \Upsilon_n = M$, then the family $\Omega_n := \Omega \cap \Upsilon_n$, $n \in \mathbb{N}$, satisfies the hypothesis of Lemma 2.8 part b).

The following definition is motivated by [13] [28], who treat the Euclidean space \mathbb{R}^m :

Definition 2.10. An open set $\Omega \subset M$ is called *Kac regular*, if one has

$$(9) \quad \mathbb{P}^x \{ \alpha_\Omega = \min(\beta_\Omega, \alpha_M) \} = 1 \quad \text{for all } x \in \Omega.$$

Note that we do not assume M to be stochastically complete, that is $\alpha_M = \infty$ \mathbb{P}^x -a.s. for all $x \in M$, noting that in the latter case one has

$$\mathbb{P}^x \{ \alpha_\Omega = \min(\beta_\Omega, \alpha_M) \} = \mathbb{P}^x \{ \alpha_\Omega = \beta_\Omega \},$$

and some technical problems are not present, that we will have to deal with. The connection between problems such as (6) and Kac regularity is clarified in the following result, which has been established in the Euclidean space \mathbb{R}^m in [13]:

Proposition 2.11. *Let $\Omega \subset M$ be an arbitrary open subset. The following properties are equivalent:*

- Ω is Kac regular.
- One has

$$(10) \quad W_0^{1,2}(\Omega) = \{ f \in W_0^{1,2}(M) : f|_{M \setminus \Omega} = 0 \text{ } \mu\text{-a.e.} \}.$$

- For all $t \geq 0$ one has

$$(11) \quad \lim_{n \rightarrow \infty} \exp(-tH_M(n1_{M \setminus \Omega})) = \exp(-tH_\Omega)P_\Omega$$

strongly as bounded operators in $L^2(M)$.

The proof of Proposition 2.11 will be given in Section 4. Note that in our notation $H_M(n1_{M \setminus \Omega})$ is the Friedrichs realization of $-\Delta + n1_{M \setminus \Omega}$ in $L^2(M)$, and H_Ω is the Dirichlet realization of $-\Delta$ in $L^2(\Omega)$. The equivalence of (10) and (11) follows immediately from specializing Proposition 2.7 to the scalar case. The equivalence of (11) and Kac-regularity will be established by using a Feynman-Kac formula.

Remark 2.12. 1. The potential theoretic variant of (10) does not see anything from the geometry of Ω , namely [4], for *every open $\Omega \subset M$* one has

$$W_0^{1,2}(\Omega) = \{ f \in W_0^{1,2}(M) : \check{f}|_{M \setminus \Omega} = 0 \text{ q.e.} \},$$

where ‘q.e.’ (quasi everywhere) is understood with respect to the capacity associated with the regular strongly local Dirichlet form

$$\langle f_1, f_2 \rangle_* = \int (df_1, df_2) d\mu$$

in $L^2(M)$ with domain of definition $W_0^{1,2}(M)$, and where \check{f} denotes the quasi-continuous representative of f . This again reflects the subtlety of the questions under investigation.

2. It was conjectured in [25] by B. Simon that open subsets $\Omega \subset \mathbb{R}^m$ of the Euclidean space with $\mathbb{R}^m \setminus \Omega$ perfect satisfy (11). A counterexample to this conjecture was given in [12] and [26] (cf. [27], p.127), who even show that there exists $\Omega \subset \mathbb{R}^m$ open with $\mathbb{R}^m \setminus \Omega$ compact and

$$\mathbb{R}^m \setminus \Omega = \overline{\mathbb{R}^m \setminus \Omega^\circ}$$

such that (10) and thus (11) fails. Note that this counterexample also entails that even for quasi-continuous f 's the assumption $f|_{M \setminus \Omega} = 0$ a.e. does not imply that $f|_{M \setminus \Omega} = 0$ q.e.

In order to formulate our main result, we add:

Definition 2.13. An open subset $\Omega \subset M$ is called *Lipschitz regular*, if there exists a family $\{\Omega_n : n \in \mathbb{N}\}$ of open subsets of Ω having Lipschitz boundary and compact closure in M , such that for all $n \in \mathbb{N}$ there exists an open subset $\Upsilon \subset M$ such that $\Upsilon_n \cap \Omega = \Omega_n$ and $\overline{\Omega} = \bigcup_{n \in \mathbb{N}} (\overline{\Omega} \cap \Upsilon_n)$.

Here comes our main result:

Theorem 2.14. *Let Ω be an arbitrary open subset of M .*

- a) *If Ω is Kac regular, then for every Kato-Schrödinger bundle $(E, \nabla, V) \rightarrow M$ one has (6) and (7).*
- b) *If Ω is Lipschitz regular, then Ω is Kac regular.*

Using transversality theory one finds that every open subset $\Omega \subset M$ having a smooth boundary is Lipschitz regular, implying:

Corollary 2.15. *Every open subset $\Omega \subset M$ having a smooth boundary is Lipschitz regular and thus Kac regular.*

Proof: The proof of the fact that Ω is Lipschitz regular follows from Lemma 18 in [24]. For the sake of completeness, we provide a sketch of proof: pick a family $\{W_n : n \in \mathbb{N}\}$ of relatively compact open subsets of M with smooth boundary such that $W_n \subset W_{n+1}$ for all $n \in \mathbb{N}$ and $\bigcup_{n \in \mathbb{N}} W_n = M$. Let $i_n : W_n \rightarrow M$ be the natural inclusion of W_n in M . Using classical results of transversality theory in differential topology we can find for each $n \in \mathbb{N}$ a smooth embedding $j_n : \overline{W}_n \rightarrow M$ such that j_n and i_n are arbitrarily close in the Whitney strong topology and $j_n(\partial \overline{W}_n)$ is transverse to $\partial \overline{\Omega}$. Altogether this tells us that $\{j_n(W_n) : n \in \mathbb{N}\}$ is a family of open subsets of M such that $j_n(W_n) \subset j(W_{n+1})$ for all $n \in \mathbb{N}$ and $\bigcup_{n \in \mathbb{N}} j(W_n) = M$ in a way that $\Omega_n := j_n(W_n) \cap \Omega$ has a Lipschitz boundary. It follows from Remark 2.9 that the set Ω becomes a Lipschitz regular set. We refer to [16] and [17] for classical results about transversality theory and for the definition of Whitney topology. ■

We **conjecture** that every open subset $\Omega \subset M$ having a locally Lipschitz boundary is Lipschitz regular.

Theorem 2.14 a) is new even in the Euclidean case, where so far only $-\Delta$ has been treated, and not even Schrödinger operators $-\Delta + V$. The proof of Theorem 2.14 will be given in Section 4. We continue with remarks on this proof: The proof of Theorem 2.14 a) relies on Proposition 2.7 and Proposition 2.11, while both worlds are linked through various partially completely new covariant Feynman-Kac formula. We believe it is a very surprising fact, that no conditions on the negative part of V is needed in Theorem 2.14 a), as such a condition is certainly required for the covariant Feynman-Kac formula to hold (roughly speaking, because Brownian paths have an infinite speed). The point here is that using rather subtle approximation arguments, one can reduce everything to the case of V 's that are bounded from below by a constant.

The proof of Theorem 2.14 b) relies again on Proposition 2.11: In a first step, we will give an analytic proof of the stronger statement

$$(12) \quad W_0^{1,2}(\Omega) = \{f \in W^{1,2}(M) : f|_{M \setminus \Omega} = 0 \text{ } \mu\text{-a.e.}\},$$

under the assumption that Ω is relatively compact with $\partial\Omega$ Lipschitz, directly verifying (10). As usual, $W^{1,2}(M)$ is defined to be the space of all $f \in L^2(M)$ such that $df \in \Gamma_{L^2}(M, T^*M)$ in the sense of distributions. In a second step we will then combine this local result with a probabilistic approximation argument, directly confirming (9).

Remark 2.16. 1. It is false that every open subset $\Omega \subset M$ with smooth boundary satisfies⁴ (12). As a counterexample, one can consider the Euclidean ball and its open upper half

$$M := \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 < 1\}, \quad \Omega := \{(x, y, z) \in M : z > 0\},$$

with the Euclidean metric.

2. It is shown in [13] that in the Euclidean case Ω 's with $\partial\Omega$ having the segment property are Kac regular. In fact, many comparable Euclidean results can be found in [26]. The segment property is more general than being locally Lipschitz, but is also a concept that does not apply to manifolds.

In view of Example 2.6.3, the following result includes (when applied to 0-forms) a criterion for a harmonic function f on an open subset $\Omega \subset M$ with $f = 0$ μ -a.e. in $M \setminus \Omega$ to have vanishing gradient in Ω (cf. [13] for the Euclidean case):

Corollary 2.17. *Let M be geodesically complete and let $(E; c, \nabla) \rightarrow M$ be a geometric Kato-Dirac bundle. Then for every open subset $\Omega \subset M$ which is Kac regular, and every Ψ with*

$$\Psi \in \Gamma_{L^2}(M, E), \quad D(c, \nabla)\Psi \in \Gamma_{L^2}(M, E), \quad \Psi = 0 \text{ } \mu\text{-a.e. in } M \setminus \Omega,$$

one has the implication

$$D(c, \nabla)^2\Psi = 0 \text{ in } \Omega \quad \Rightarrow \quad D(c, \nabla)\Psi = 0 \text{ in } \Omega,$$

⁴But of course this statement is true by Proposition 2.11 and Theorem 2.14 b), if M is geodesically complete, as then $W^{1,2}(M) = W_0^{1,2}(M)$ [1].

where above the action of $D(c, \nabla)^2$ and $D(c, \nabla)$ is understood in the distributional sense.

Proof: Recall that $(E; c, \nabla) \rightarrow M$ induces the Kato-Schrödinger bundle

$$(E, \nabla, V(c, \nabla)) \longrightarrow M,$$

As M is geodesically complete one has

$$(13) \quad \Psi \in \Gamma_{W_0^{1,2}}(M, E; \nabla, V(c, \nabla)),$$

which in view of the product rule for $D(c, \nabla)$ (p. 171 in [11]) and a manifold version of the Meyers-Serrin Theorem (Theorem 2.9 in [15]) can be seen as in the case of functions [1]. Now (13) and the Kac regularity of Ω imply

$$\Psi \in \Gamma_{W_0^{1,2}}(\Omega, E, \nabla, V(c, \nabla)),$$

and we can pick a sequence $\Psi_n \in \Gamma_{C^\infty}(\Omega, E)$ with

$$\lim_{n \rightarrow \infty} \|\Psi_n - \Psi\|_{\nabla, V(c, \nabla)} = 0.$$

It follows that

$$\begin{aligned} \|D(c, \nabla)\Psi\|^2 &= \lim_{n \rightarrow \infty} \langle D(c, \nabla)\Psi_n, D(c, \nabla)\Psi \rangle \\ &= \int_{\Omega} (D(c, \nabla)\Psi_n, D(c, \nabla)\Psi) d\mu = \int_{\Omega} (\Psi_n, D(c, \nabla)^2\Psi) d\mu = 0, \end{aligned}$$

where the integration by parts is justified as Ψ is smooth in Ω by local elliptic regularity, and Ψ_n is smooth and compactly supported in Ω . \blacksquare

Note that in Corollary 2.17 the geodesic completeness of M was only assumed to guarantee

$$\Psi \in \Gamma_{W_0^{1,2}}(M, E; \nabla, V(c, \nabla)),$$

which one could assume instead.

3. Proof of Lemma 2.8

Proof of Lemma 2.8: a) This statement is well-known and elementary to check.

b) As this observation seems to be new, we have decided to add its proof (which is rather technical): Assume first that there exists $t_0 \in (0, \infty)$ such that $\beta_{\Omega}(\omega) = t_0$. This means that the Lebesgue measure of the set $\{s \in [0, t_0] : \omega(s) \notin \Omega\}$ is zero, briefly

$$|\{s \in [0, t_0] : \omega(s) \notin \Omega\}| = 0$$

and moreover that for each $\epsilon > 0$ we have

$$|\{s \in [0, t_0 + \epsilon] : \omega(s) \notin \Omega\}| > 0.$$

Let $n_0 > 0$ be such that

$$\omega([0, t_0]) \cap \Omega = \omega([0, t_0]) \cap \Omega_{n_0}.$$

In this way we have

$$|\{s \in [0, t_0] : \omega(s) \notin \Omega_n\}| = 0$$

for each $n \geq n_0$, because

$$\{s \in [0, t_0] : \omega(s) \notin \Omega_n\} = \{s \in [0, t_0] : \omega(s) \notin \Omega_{n_0}\} = \{s \in [0, t_0] : \omega(s) \notin \Omega\}$$

and the latter has Lebesgue measure 0. Moreover we have

$$|\{s \in [0, t_0 + \delta] : \omega(s) \notin \Omega_n\}| > 0$$

for each $\delta > 0$ and for each $n \geq n_0$, because

$$\{s \in [0, t_0 + \delta] : \omega(s) \notin \Omega\} \subset \{s \in [0, t_0 + \delta] : \omega(s) \notin \Omega_n\} \subset \{s \in [0, t_0 + \delta] : \omega(s) \notin \Omega_{n_0}\}$$

and

$$|\{s \in [0, t_0 + \delta] : \omega(s) \notin \Omega\}| > 0.$$

Therefore $\beta_{\Omega_n}(\omega) = t_0$ for each $n \geq n_0$ and hence we can conclude that $\lim \beta_{\Omega_n}(\omega) = \beta_{\Omega}(\omega)$ as $n \rightarrow \infty$.

Assume now that there is no $t_0 \in (0, \infty)$ such that $\beta_{\Omega}(\omega) = t_0$. Therefore

$$|\{s \in [0, \infty) : \omega(s) \notin \Omega\}| = 0.$$

In this case we set $\beta_{\Omega}(\omega) := \infty$. Consider first the case where $\overline{\omega([0, \infty))}$ is compact. Let

$$A := \{s \in [0, \infty) : \omega(s) \notin \Omega\}$$

and let $B := [0, \infty) \setminus A$. Let n_0 be such that $\omega(B) \subset \Omega_{n_0}$. Since $\omega(B) \subset \Omega_{n_0}$ we have

$$\omega([0, \infty)) \cap \Omega = \omega([0, \infty)) \cap \Omega_n$$

for each $n \geq n_0$. Therefore

$$|\{s \in [0, \infty) : \omega(s) \notin \Omega_n\}| = 0$$

for any $n \geq n_0$ and hence we can conclude that $\lim \beta_{\Omega_n}(\omega) = \beta_{\Omega}(\omega) = \infty$ as $n \rightarrow \infty$. Finally consider the case where $\overline{\omega([0, \infty))}$ is not compact. The sequence $\beta_{\Omega_n}(\omega)$ is increasing and unbounded, because given an arbitrarily big t_0 we can find an integer n_0 such that $\omega([0, t_0]) \cap \Omega = \omega([0, t_0]) \cap \Omega_{n_0}$ and $\omega(t_1) \notin \Omega_{n_0}$ for some $t_1 > t_0$. Hence $\beta_{\Omega_{n_0}}(\omega) \geq t_0$ and therefore $\lim \beta_{\Omega_n}(\omega) = \infty$ as $n \rightarrow \infty$. \blacksquare

4. Proofs of main results

We first introduce some notation that will be relevant in the sequel. Let

$$(E, V, \nabla) \longrightarrow M$$

be a Kato-Schrödinger bundle.

Notation 4.1. Given an open subset $\Omega \subset M$ let

(14)

$$\Gamma_{\widetilde{W}_0^{1,2}}(\Omega, E; \nabla, V) := \{f \in \Gamma_{W_0^{1,2}}(M, E; \nabla, V) : f|_{M \setminus \Omega} = 0 \text{ } \mu\text{-a.e.}\} \subset \Gamma_{W_0^{1,2}}(M, E; \nabla, V),$$

and let $\widetilde{H}_{\Omega}(\nabla, V)$ denote the self-adjoint nonnegative operator in $\Gamma_{L^2}(\Omega, E)$ which corresponds to $\langle \cdot, \cdot \rangle_{\nabla, V, *}$ with domain of definition $\Gamma_{\widetilde{W}_0^{1,2}}(\Omega, E; \nabla, V)$, a closed densely defined

symmetric sesquilinear form in $\Gamma_{L^2}(\Omega, E)$. Again we will follow the conventions from (2.4) analogously.

The above auxiliary operator $\tilde{H}_\Omega(\nabla, V)$ satisfies the following two important results, without any further assumptions on Ω :

Proposition 4.2. *For every open subset Ω of M and every $t \geq 0$ one has*

$$\lim_{n \rightarrow \infty} \exp(-tH_M(\nabla, V + n1_{M \setminus \Omega})) = \exp(-t\tilde{H}_\Omega(\nabla, V))P_\Omega$$

strongly as bounded operators in $\Gamma_{L^2}(M, E)$.

Proof: This is a simple application of Theorem 5.1: Regarding $n1_{M \setminus \Omega}$ as a bounded multiplication operator it follows that with

$$Q_n := \langle \cdot, \cdot \rangle_{\nabla, V + n1_{M \setminus \Omega}, * } = \langle \cdot, \cdot \rangle_{\nabla, V, * } + n \int_{M \setminus \Omega} (\cdot, \cdot) d\mu, \quad \text{Dom}(Q_n) := \Gamma_{W_0^{1,2}}(M, E; \nabla, V),$$

one has

$$S_{Q_n} = H_M(\nabla, V + n1_{M \setminus \Omega}).$$

It follows that $f \in \Gamma_{W_0^{1,2}}(M, E; \nabla, V)$ is in $\text{Dom}(Q_\infty)$, if and only if

$$\sup_{n \in \mathbb{N}} n \int_{M \setminus \Omega} |f|^2 d\mu < \infty,$$

which again is equivalent to $f = 0$ in $M \setminus \Omega$ μ -a.e., and for such f 's one has

$$Q_\infty(f, f) = \lim_{n \rightarrow \infty} Q_n(f, f) = \langle f, f \rangle_{\nabla, V, * },$$

proving $S_{Q_\infty} = \tilde{H}_\Omega(\nabla, V)$. ■

Let us now prepare the formulation of various covariant Feynman-Kac formulae, that is, path integral representations of the semigroups corresponding to covariant Schrödinger operators. Note that the usual Feynman-Kac formula [9, 18] is for semigroups of the form $\exp(-tH_M(\nabla, V))$, and it is the aim of this section to derive formulae for the semigroups $\exp(-t\tilde{H}_\Omega(\nabla, V))$ and $\exp(-tH_\Omega(\nabla, V))$. As these formulae rely on semimartingale theory and stochastic integrals (through the definition of the Stratonovic parallel transport), one needs to work here with a filtered probability, which allows to pick continuous versions of these semimartingales. We refer the reader to [18, 9] for the basics of stochastic analysis on manifolds. To set the stage, suppose that for every $x \in M$ we are given an adapted Riemannian Brownian motion $\mathbb{X}(x)$ on M with

$$\mathbb{P}\{\mathbb{X}_0(x) = x\} = 1,$$

which is defined on some fixed filtered probability space which satisfies the usual assumptions of completeness and right-continuity. In other words, $\mathbb{X}(x)$ is an M -valued continuous and adapted process defined up to its explosion time, such that the law of $\mathbb{X}(x)$ is \mathbb{P}^x . Given an open subset $\Omega \subset M$, let $\alpha_\Omega(x) := \alpha_\Omega(\mathbb{X}(x))$ denote the first exit time of $\mathbb{X}(x)$ from Ω , and let $\beta_\Omega(x) := \beta_\Omega(\mathbb{X}(x))$ denote the penetration time of $\mathbb{X}(x)$ to $M \setminus \Omega$. Let $//_t^\nabla(x)$,

$t < \alpha_M(x)$, be the Stratonovic parallel transport with respect to ∇ along the paths of $\mathbb{X}(x)$, which, ∇ being metric, is a random variable taking values in the unitary maps $E_x \rightarrow E_{\mathbb{X}(t)}$. So far $x \in M$ was arbitrary. On the other hand, for μ -a.e. $x \in M$, we can define pathwise for $t < \alpha_M(x)$ a random variable $\mathcal{A}_V^\nabla(x, t)$ taking values in $\text{End}(E_x)$, to be the unique locally absolutely continuous solution of

$$\begin{aligned} (d/dt)\mathcal{A}_V^\nabla(x, t) &= -\mathcal{A}_V^\nabla(x, t)(//_t^\nabla(x)^{-1}V(\mathbb{X}_t(x))//_t^\nabla(x)), \\ \mathcal{A}_V^\nabla(x, 0) &= 1_{\text{End}(E_x)}. \end{aligned}$$

The existence and uniqueness of $\mathcal{A}_V^\nabla(x, t)$ is guaranteed by the L_{loc}^1 -assumption on V , which implies

$$\mathbb{P} \{ |V(\mathbb{X}(x))| \in L_{\text{loc}}^1[0, \alpha_M(x)] \} = 1 \quad \text{for } \mu\text{-a.e. } x \in M,$$

and so

$$\mathbb{P} \{ |//_t^\nabla(x)^{-1}V(\mathbb{X}_t(x))//_t^\nabla(x)| \in L_{\text{loc}}^1[0, \alpha_M(x)] \} = 1 \quad \text{for } \mu\text{-a.e. } x \in M.$$

We refer the reader to [9] for detailed proofs of these facts.

Proposition 4.3. *For every $\Omega \subset M$ open, $t \geq 0$, $f \in \Gamma_{L^2}(\Omega, E)$, and μ -a.e. $x \in M$ one has*

$$(15) \quad \exp(-t\tilde{H}_\Omega(\nabla, V))f(x) = \int_{\{t < \min(\beta_\Omega(x), \alpha_M(x))\}} \mathcal{A}_V^\nabla(x, t)//_t^\nabla(x)^{-1}f(\mathbb{X}_t(x))d\mathbb{P}.$$

Note that the expectation in (15) is on the fiber E_x , as $f(\mathbb{X}_t(x)) \in E_{\mathbb{X}_t(x)}$ in $\{t < \alpha_M(x)\}$. In the scalar case we get

$$(16) \quad \exp(-t\tilde{H}_\Omega(w))f(x) = \int_{\{t < \min(\beta_\Omega(x), \alpha_M(x))\}} \exp(-\int_0^t w(\mathbb{X}_s))f(\mathbb{X}_t)d\mathbb{P}.$$

Proof of Proposition 4.3: We start by recalling the usual covariant Feynman-Kac formula [9]

$$\exp(-tH_M(\nabla, W))f(x) = \int_{\{t < \alpha_M(x)\}} \mathcal{A}_W^\nabla(x, t)//_t^\nabla(x)^{-1}f(\mathbb{X}_t(x))d\mathbb{P},$$

which is valid, for example if the potential W is in L_{loc}^2 is such that for some constant $A \in \mathbb{R}$ and all $x \in M$, the eigenvalues of $W(x) \in \text{End}(E_x)$ are $\geq A$. In particular, this statement includes that

$$(17) \quad \int_{\{t < \alpha_M(x)\}} |\mathcal{A}_W^\nabla(x, t)//_t^\nabla(x)^{-1}f(\mathbb{X}_t(x))| d\mathbb{P} < \infty.$$

Applying this with $W = V + n1_{M \setminus \Omega}$ and using Proposition 4.2 we have, possibly after taking a subsequence of

$$\exp(-tH_M(\nabla, V + n1_{M \setminus \Omega}))f$$

if necessary, that for μ -a.e. $x \in M$ it holds that

$$\begin{aligned} \exp(-t\tilde{H}_\Omega(\nabla, V))f(x) &= \lim_{n \rightarrow \infty} \exp(-tH_M(\nabla, V + n1_{M \setminus \Omega}))f(x) \\ &= \lim_{n \rightarrow \infty} \int_{\{t < \alpha_M(x)\}} \exp\left(-n \int_0^t 1_{M \setminus \Omega}(\mathbb{X}_s(x))ds\right) \mathcal{A}_V^\nabla(x, t) //_t^\nabla(x)^{-1} f(\mathbb{X}_t(x)) d\mathbb{P}. \end{aligned}$$

For paths from the set

$$\{t < \alpha_M(x)\} =_{\mathbb{P}} \{\mathbb{X}_s(x) \in M \text{ for all } s \in [0, t]\}$$

one has

$$\int_0^t 1_{\hat{M} \setminus \Omega}(\mathbb{X}_s(x))ds = \int_0^t 1_{M \setminus \Omega}(\mathbb{X}_s(x))ds,$$

and $\int_0^t 1_{M \setminus \Omega}(\mathbb{X}_s(x))ds = 0$ is equivalent to $t < \beta_\Omega(x)$. Thus, \mathbb{P} -a.s. in $\{t < \alpha_M(x)\}$ one has

$$\lim_{n \rightarrow \infty} \exp\left(-n \int_0^t 1_{M \setminus \Omega}(\mathbb{X}_s(x))ds\right) = 1_{\{t < \beta_\Omega(x)\}}$$

and

$$\begin{aligned} &\lim_{n \rightarrow \infty} \int_{\{t < \alpha_M(x)\}} \exp\left(-n \int_0^t 1_{M \setminus \Omega}(\mathbb{X}_s(x))ds\right) \mathcal{A}_V^\nabla(x, t) //_t^\nabla(x)^{-1} f(\mathbb{X}_t(x)) d\mathbb{P} \\ &= \int_{\{t < \min(\beta_\Omega(x), \alpha_M(x))\}} \mathcal{A}_V^\nabla(x, t) //_t^\nabla(x)^{-1} f(\mathbb{X}_t(x)) d\mathbb{P} \end{aligned}$$

follows from dominated convergence, in view of (17). ■

4.1. Proof of Proposition 2.7.

Proof of Proposition 2.7: With the preparations made, there is almost nothing left to prove: Note first that (6) is equivalent to

$$(18) \quad H_\Omega(\nabla, V) = \tilde{H}_\Omega(\nabla, V).$$

(6) \Rightarrow (7): We have (18), so that (7) follows from Proposition 4.2.

(7) \Rightarrow (6): (7) and Proposition 4.2 imply

$$\exp(-tH_\Omega(\nabla, V)) = \exp(-t\tilde{H}_\Omega(\nabla, V)) \quad \text{for all } t > 0,$$

showing (18). ■

4.2. Proof of Proposition 2.11.

Proof of Proposition 2.11: Note first that one has the Feynman-Kac formula

$$\exp(-tH_\Omega)f(x) = \int_{\{t < \alpha_\Omega\}} f(\mathbb{X}_t) d\mathbb{P}^x$$

for all $f \in L^2(\Omega)$, $t > 0$, $x \in \Omega$, meaning also that the right hand side is the smooth representative of $\exp(-tH_\Omega)f$. In case Ω is relatively compact with smooth boundary this is well-known. For the general case we can assume $f \geq 0$. Fix $x \in \Omega$ and exhaust Ω with a sequence of relatively compact smooth Ω_n 's with $x \in \Omega_1$. Then the latter formula is valid with Ω replaced with Ω_n , and we can take $n \rightarrow \infty$ using monotone convergence for integrals on the probabilistic side, noting that $\alpha_{\Omega_n} \nearrow \alpha_\Omega$ by Lemma 2.8 a), and the fact that (cf. p.213 in [7])

$$\exp(-tH_{\Omega_n})f(x) \nearrow \exp(-tH_\Omega)f(x).$$

Let us come to the actual proof of Proposition 2.11. Being equipped with the previously established results, we can follow the Euclidean proof from [13] from here on:

As we have already remarked, the equivalence of (10) and (11) follows immediately from Proposition 2.7.

(10) \Rightarrow *Kac-regularity:* (10) implies

$$(19) \quad \tilde{H}_\Omega = H_\Omega,$$

by comparing the associated forms. In particular, the semigroups coincide and the Feynman-Kac formulae for the semigroups imply the first identity in

$$0 = \int (1_{\{t < \min(\beta_\Omega, \alpha_M)\}} - 1_{\{t < \alpha_\Omega\}}) f(\mathbb{X}_t) d\mathbb{P}^x = \int_{\{\alpha_\Omega \leq t < \beta_\Omega\}} f(\mathbb{X}_t) d\mathbb{P}^x$$

for all $f \in L^2(\Omega)$, $t > 0$, μ -a.e. $x \in \Omega$, where the second equality follows from

$$1_{\{\alpha_\Omega \leq t < \min(\beta_\Omega, \alpha_M)\}} = 1_{\{t < \min(\beta_\Omega, \alpha_M)\}} - 1_{\{t < \alpha_\Omega\} \cap \{t < \min(\beta_\Omega, \alpha_M)\}}$$

and $\alpha_\Omega \leq \beta_\Omega$ and $\alpha_\Omega \leq \alpha_M$. Letting f run through $f_n := 1_{A_n}$, where $\{A_n : n \in \mathbb{N}\}$ is a family of compact subsets of M such that $A_n \subset A_{n+1}$ for all $n \in \mathbb{N}$ and $\bigcup_{n \in \mathbb{N}} A_n = M$, we get

$$\mathbb{P}^x \{\alpha_\Omega \leq t < \min(\beta_\Omega, \alpha_M)\} = 0 \quad \text{for all } t > 0, \mu\text{-a.e. } x \in \Omega.$$

Letting t run through $\mathbb{Q}_{>0}$,

$$(20) \quad \mathbb{P}^x \{\alpha_\Omega < \min(\beta_\Omega, \alpha_M)\} = 0 \quad \text{for } \mu\text{-a.e. } x \in \Omega.$$

In view of $\alpha_\Omega \leq \beta_\Omega$, the proof of the Kac-regularity of Ω is complete, once we have shown that the bounded function $h(x) := \mathbb{P}^x \{\alpha_\Omega < \min(\beta_\Omega, \alpha_M)\}$ is continuous in Ω . We are going to prove that for all open relatively compact subsets $D \subset \Omega$ with $\bar{D} \subset \Omega$ one has

$$(21) \quad \int h(\mathbb{X}_{\alpha_D}) d\mathbb{P}^x = h(x) \quad \text{for all } x \in D,$$

which implies that $x \mapsto \mathbb{P}^x \{\alpha_\Omega < \min(\beta_\Omega, \alpha_M)\}$ is harmonic and thus smooth in Ω . To see (21), note first that $\alpha_D < \alpha_\Omega \leq \alpha_M$ \mathbb{P}^x -a.s., as the transition density of Brownian motion is continuous. Now we can calculate

$$\begin{aligned}
& \int h(\mathbb{X}_{\alpha_D}) d\mathbb{P}^x \\
&= \int \mathbb{P}^{\mathbb{X}_{\alpha_D}} \{\alpha_\Omega < \min(\beta_\Omega, \alpha_M)\} d\mathbb{P}^x \\
&= \int 1_{\{\alpha_\Omega < \min(\beta_\Omega, \alpha_M)\}} (\omega(\alpha_D(\omega) + \bullet)) d\mathbb{P}^x(\omega) \\
&= \int 1_{\{\alpha_\Omega + \alpha_D < \min(\beta_\Omega, \alpha_M) + \alpha_D\}} (\omega) d\mathbb{P}^x(\omega) \\
&= \int 1_{\{\alpha_\Omega < \min(\beta_\Omega, \alpha_M)\}} (\omega) d\mathbb{P}^x(\omega) \\
&= h(x),
\end{aligned}$$

where the first inequality holds by definition, the second equality holds by the strong Markov property of Brownian motion, the third equality holds by elementary considerations, the fourth equality is trivial and the last equality again holds by definition.

Kac-regularity \Rightarrow (10): By the corresponding Feynman-Kac formulae we immediately find that Kac-regularity implies

$$\exp(-t\tilde{H}_\Omega) = \exp(-tH_\Omega) \quad \text{for all } t > 0,$$

so that $\tilde{H}_\Omega = H_\Omega$. ■

4.3. *Proof of Theorem 2.14 a).* We will need the following covariant-Feynman-Kac formula:

Proposition 4.4. *For every $\Omega \subset M$ open, $t \geq 0$, $f \in \Gamma_{L^2}(\Omega, E)$, and μ -a.e. $x \in M$ one has the covariant Feynman-Kac formula*

$$(22) \quad \exp(-tH_\Omega(\nabla, V))f(x) = \int_{\{t < \alpha_\Omega(x)\}} \mathcal{A}_V^\nabla(x, t) //_t^\nabla(x)^{-1} f(\mathbb{X}_t(x)) d\mathbb{P}.$$

In case Ω is relatively compact with smooth boundary and V is smooth, formula (22) is implicitly included in [3]. For general Ω 's and V 's (22) seems to be new.

Proof of Proposition 4.4: It is enough to prove the formula in case Ω is relatively compact with a smooth boundary. Indeed, in the general case we can then pick a family of open subsets $\{\Omega_n : n \in \mathbb{N}\}$ of Ω having a smooth boundary, such that $\Omega_n \subset \Omega_{n+1}$ for all $n \in \mathbb{N}$ and $\bigcup_{n \in \mathbb{N}} \Omega_n = \Omega$. Then we have $\alpha_{\Omega_n}(x) \nearrow \alpha_\Omega(x)$ and

$$(23) \quad \int_{\{t < \alpha_{\Omega_n}(x)\}} \mathcal{A}_V^\nabla(x, t) //_t^\nabla(x)^{-1} f(\mathbb{X}_t(x)) d\mathbb{P} \rightarrow \int_{\{t < \alpha_\Omega(x)\}} \mathcal{A}_V^\nabla(x, t) //_t^\nabla(x)^{-1} f(\mathbb{X}_t(x)) d\mathbb{P}.$$

follows from dominated convergence. Likewise,

$$(24) \quad \exp(-tH_{\Omega_n}(\nabla, V))f(x) \rightarrow \exp(-tH_{\Omega}(\nabla, V))f(x)$$

follows from applying Theorem 5.2 and Remark 5.3 thereafter: indeed, define the form Q by

$$\text{Dom}(Q) := \Gamma_{W_0^{1,2}}(\Omega, E; \nabla, V), \quad Q(f_1, f_2) = \langle f_1, f_2 \rangle_{\nabla, V, *}, \quad f_1, f_2 \in \text{Dom}(Q).$$

Then the form Q' from Theorem 5.2 is closable (it has the closed extension Q), so that it remains to prove $\overline{Q'} = Q$, which is easily checked from the definitions.

Thus we can and we will assume Ω is relatively compact and has a smooth boundary. In case V is smooth and bounded, the asserted formula is a simple consequence of Ito's formula and parabolic regularity up to the boundary (cf. p. 102 in [3]). In case V is bounded, we can use Friedrichs mollifiers to pick a sequence of smooth potentials $V_n : M \rightarrow \text{End}(E)$ with $|V_n| \leq |V|$ and $|V_n - V| \rightarrow 0$ μ -a.e. in M . Applying (22) with V replaced with V_n and taking $n \rightarrow \infty$ gives the result in this case. Finally, if V is bounded from below, we can use the spectral calculus on the fibers of $E \rightarrow M$ to define a sequence of potentials

$$V_n : M \rightarrow \text{End}(E), \quad V_n(x) := \min(n, V(x))$$

then apply (22) with V replaced with V_n and take $n \rightarrow \infty$. Each approximation argument can be justified precisely as in the proof of Theorem 2.11 in [9], which treats the case $M = \Omega$: In each situation one uses convergence results for sesquilinear forms to control the left-hand side of (22) and convergence theorems for integrals to control the right-hand-side. \blacksquare

Proof of Theorem 2.14 a): There is nothing left to prove: We know from Proposition 4.2 that

$$\lim_{n \rightarrow \infty} \exp(-tH_M(\nabla, V + n1_{M \setminus \Omega})) = \exp(-t\widetilde{H}_{\Omega}(\nabla, V))P_{\Omega}$$

strongly (without any further assumptions on Ω), and

$$\widetilde{H}_{\Omega}(\nabla, V) = H_{\Omega}(\nabla, V)$$

follows from a comparison of the covariant Feynman-Kac formulae for $\exp(-t\widetilde{H}_{\Omega}(\nabla, V_n))$ and $\exp(-tH_{\Omega}(\nabla, V_n))$ (cf. Proposition 4.3 and Proposition 4.4), using that Ω is Kac regular. \blacksquare

4.4. *Proof of Theorem 2.14 b).* We recall that given an open subset $\Omega \subset M$, the space $W^{1,2}(\Omega; d)$ is defined to be the space of all $f \in L^2(\Omega)$ such that $df \in \Gamma_{L^2}(\Omega; T^*M)$ in the sense of distributions. We are going to need a special case of the following result for the proof of Theorem 2.14 b):

Proposition 4.5. *Let $\Omega \subset M$ be a relatively compact open subset with a Lipschitz boundary. Then one has*

$$W_0^{1,2}(\Omega) = \{f \in W^{1,2}(M) : f|_{M \setminus \Omega} = 0 \text{ } \mu\text{-a.e.}\},$$

in particular, Ω is Kac-regular.

Proof: Set $m := \dim M$. It remains to prove

$$W_0^{1,2}(\Omega) \supset \{f \in W^{1,2}(M) : f|_{M \setminus \Omega} = 0 \text{ } \mu\text{-a.e.}\}.$$

To this end, we record that for every $p \in \overline{\Omega} \setminus \Omega = \partial\Omega$ there exists a pair of open neighborhoods V, X of p , and a smooth diffeomorphism $\alpha : X \rightarrow A \subset \mathbb{R}^m$ such that:

- (1) $\overline{V} \subset X$,
- (2) \overline{V} is compact,
- (3) $\alpha(V \cap \Omega)$ is a bounded Lipschitz set of \mathbb{R}^m .

Let us consider a finite collection of open subsets of M , $\mathcal{W} := \{V_1, \dots, V_q, W_1, \dots, W_n\}$, such that

- for each $j \in \{1, \dots, n\}$ we have $\overline{W_j} \subset \Omega$,
- $\overline{\Omega} \subset V_1 \cup \dots \cup V_q \cup W_1 \cup \dots \cup W_n$,
- for each $i \in \{1, \dots, q\}$ there exists another open subset X_i such that the pair given by V_i and X_i satisfies the properties (1)–(3) required in the statement.

Clearly such a finite collection of open subsets exists. Let I be an open subset of M such that $I \cap \Omega = \emptyset$ and

$$M = V_1 \cup \dots \cup V_q \cup W_1 \cup \dots \cup W_n \cup I.$$

Let

$$\mathcal{W}' := \{V_1, \dots, V_q, W_1, \dots, W_n, I\}$$

and let

$$\{\phi_1, \dots, \phi_q, \psi_1, \dots, \psi_n, \tau\}$$

be a partition of unity subordinated to \mathcal{W} . Given now $f \in W^{1,2}(M)$ with $f|_{M \setminus \Omega} = 0$ μ -a.e., for each $i \in \{1, \dots, n\}$ we have $\text{supp}(\psi_i f) \subset W_i$. Therefore there exists a sequence $\{\beta_p^i\}_{p \in \mathbb{N}} \subset C_c^\infty(W_i)$ such that $\beta_p^i \rightarrow \psi_i f$ in $W_0^{1,2}(W_i)$ as $p \rightarrow \infty$. Indeed, by a Meyers-Serrin type theorem [15], we can pick a sequence $\{\overline{\beta}_p^i\}_{p \in \mathbb{N}}$ in $C^\infty(W_i) \cap W^{1,2}(W_i)$ such that $\overline{\beta}_p^i \rightarrow f$ in $W^{1,2}(W_i)$. Then, by defining $\beta_p^i := \psi_i \overline{\beta}_p^i$, it is clear that $\{\beta_p^i\}_{p \in \mathbb{N}} \subset C_c^\infty(W_i)$ and that $\beta_p^i \rightarrow \psi_i f$ in $W_0^{1,2}(W_i)$ as $p \rightarrow \infty$. Let us now consider the other case. For each $i \in \{1, \dots, q\}$ let $Y_i := V_i \cap \Omega$. Then we have $\text{supp}(\phi_i f) \subset (\overline{Y_i} \cap V_i)$. Let $A_i := \alpha_i(X_i)$, $E_i := \alpha_i(V_i)$ and $B_i := \alpha_i(Y_i)$. Since we assumed that $\overline{V_i} \subset X_i$ we have that $(\alpha_i^* g_e)|_{V_i}$ is quasi-isometric to $g|_{V_i}$ where g_e is the standard Euclidean metric on \mathbb{R}^m and g the metric on M . In this way we can conclude that $(\phi_i f) \circ (\alpha|_{V_i})^{-1} \in W_{0,e}^{1,2}(E_i)$ and that $(\phi_i f) \circ (\alpha|_{Y_i})^{-1} \in W_e^{1,2}(B_i)$, where of course $W_e^{1,2}$ stands for the various Sobolev spaces that are defined with respect to g_e . Furthermore we know that $(\phi_i f) \circ (\alpha|_{Y_i})^{-1}|_{A_i \setminus B_i} \equiv 0$ because $(\phi_i f)|_{X_i \setminus Y_i} \equiv 0$. Therefore, by extending $(\phi_i f) \circ (\alpha|_{Y_i})^{-1}$ outside its support as the identically zero function, we can say, with a little abuse of notation, that $(\phi_i f) \circ (\alpha|_{Y_i})^{-1} \in W_e^{1,2}(\mathbb{R}^m)$ and that $(\phi_i f) \circ (\alpha|_{Y_i})^{-1}|_{\mathbb{R}^m \setminus B_i} \equiv 0$. As B_i is a bounded Lipschitz set in \mathbb{R}^m , it follows from Remark 2.16.2 that there exists a sequence $\{v_p^i\}_{p \in \mathbb{N}} \subset C_c^\infty(B_i)$ such that $v_p^i \rightarrow (\phi_i f) \circ (\alpha|_{Y_i})^{-1}$ in $W_{0,e}^{1,2}(B_i)$ as $p \rightarrow \infty$. Finally, using the fact that $(\alpha_i^* g_e)|_{V_i}$ is quasi-isometric to $g|_{V_i}$, we can conclude that $\gamma_p^i \rightarrow \phi_i f$ in $W_0^{1,2}(Y_i)$ as $p \rightarrow \infty$ where $\gamma_p^i := v_p^i \circ (\alpha_i|_{Y_i})$ and clearly $\{\gamma_p^i\}_{p \in \mathbb{N}} \subset C_c^\infty(Y_i)$ by construction. Let us define now the following sequence of functions

$\eta_p := \gamma_p^1 + \dots + \gamma_p^q + \beta_p^1 + \dots + \beta_p^n$. It is clear by construction that $\{\eta_p\}_{p \in \mathbb{N}} \subset C_c^\infty(\Omega)$. Moreover we have

$$\begin{aligned} \|\eta_p - f\| &= \|\gamma_p^1 + \dots + \gamma_p^q + \beta_p^1 + \dots + \beta_p^n - \sum_{i=1}^q \phi_i f - \sum_{i=1}^n \psi_i f\| \\ &\leq \|\gamma_p^1 - \phi_1 f\| + \dots + \|\gamma_p^q - \phi_q f\| + \|\beta_p^1 - \psi_1 f\| + \dots + \|\beta_p^n - \psi_n f\| \end{aligned}$$

and by construction all the terms in the second line tend to zero as $p \rightarrow \infty$. Hence we have shown that $\eta_j \rightarrow f$ in $\Gamma_{L^2}(\Omega, T_{\mathbb{C}}^*M)$ as $p \rightarrow \infty$. Similarly we have

$$\begin{aligned} \|d\eta_p - df\| &= \|d\gamma_p^1 + \dots + d\gamma_p^q + d\beta_p^1 + \dots + d\beta_p^n - d(\sum_{i=1}^q \phi_i f - \sum_{i=1}^n \psi_i f)\| \\ &\leq \|d\gamma_p^1 - d(\phi_1 f)\| + \dots + \|d\gamma_p^q - d(\phi_q f)\| + \|d\beta_p^1 - d(\psi_1 f)\| + \dots \\ &\quad + \|d\beta_p^n - d(\psi_n f)\| \end{aligned}$$

and again by construction we know that all the terms on the right hand side of the inequality tend to zero as $p \rightarrow \infty$. This tells us that $d\eta_p \rightarrow df$ in $\Gamma_{L^2}(\Omega, T_{\mathbb{C}}^*M)$. In conclusion we have shown that $\eta_p \rightarrow f$ in $W^{1,2}(\Omega)$ and thereby we can conclude that $f \in W_0^{1,2}(\Omega)$ as desired. \blacksquare

Proof of Theorem 2.14 b): Fix $x \in \Omega$. Pick a sequence $\Omega_n \subset \Omega$, $n \in \mathbb{N}$ of relatively compact open subsets of M with Lipschitz boundary, such that $x \in \Omega_1$, $\Omega_n \subset \Omega_{n+1}$ for all $n \in \mathbb{N}$, $\bigcup_{n \in \mathbb{N}} \Omega_n = \Omega$ and such that for all $n \in \mathbb{N}$ there exists an open subset $\Upsilon \subset M$ such that $\Upsilon_n \cap \Omega = \Omega_n$ for all $n \in \mathbb{N}$ and $\bigcup_{n \in \mathbb{N}} (\Upsilon_n \cap \bar{\Omega}) = \bar{\Omega}$. Then by Proposition 2.14 and Proposition 4.5 we have

$$\mathbb{P}^x \{\alpha_{\Omega_n} = \min(\beta_{\Omega_n}, \alpha_M)\} = 1 \quad \text{for all } n,$$

so that using $\alpha_{\Omega_n} \rightarrow \alpha_\Omega$ and $\beta_{\Omega_n} \rightarrow \beta_\Omega$ \mathbb{P}^x -a.s. as $n \rightarrow \infty$, a consequence of Lemma 2.8, we arrive at

$$\mathbb{P}^x \{\alpha_\Omega \neq \min(\beta_\Omega, \alpha_M)\} \subset \mathbb{P}^x \bigcup_{n \in \mathbb{N}} \{\alpha_{\Omega_n} \neq \min(\beta_{\Omega_n}, \alpha_M)\} \leq \sum_{n \in \mathbb{N}} \mathbb{P}^x \{\alpha_{\Omega_n} \neq \min(\beta_{\Omega_n}, \alpha_M)\} = 0,$$

completing the proof. \blacksquare

5. Appendix: Some functional analytic facts

In this section we collect some facts about the monotone convergence of nonnegative closed sesquilinear forms which are possibly not densely defined.

Let \mathcal{H} be a complex separable Hilbert space and assume that Q is a semibounded from below closed sesquilinear forms such a form on \mathcal{H} . Then Q is a densely defined semibounded from below closed sesquilinear form on the Hilbert subspace $\mathcal{H}_Q := \overline{\text{Dom}(Q)}^{\|\cdot\|_{\mathcal{H}}} \subset \mathcal{H}$

and thus there exists a unique semibounded from below self-adjoint operator S_Q on \mathcal{H}_Q such that

$$\text{Dom}(\sqrt{S_Q}) = \text{Dom}(Q), \quad Q(f_1, f_2) = \left\langle \sqrt{S_Q}f_1, \sqrt{S_Q}f_2 \right\rangle_{\mathcal{H}}.$$

Let then $P_Q : \mathcal{H} \rightarrow \mathcal{H}_Q$ denote the orthogonal projection. Recall also that for semi-bounded forms Q_1 and Q_2 in \mathcal{H} one per definitionem has $Q_1 \leq Q_2$ if and only if $\text{Dom}(Q_1) \supset \text{Dom}(Q_2)$ and $Q_1(f, f) \leq Q_2(f, f)$ for all $f \in \text{Dom}(Q_2)$. Given a semi-bounded sesquilinear form Q on \mathcal{H} there exists a largest (with respect to ' \leq ') closable semibounded sesquilinear form $\text{ref}(Q)$ on \mathcal{H} , such that $\text{ref}(Q) \leq Q$.

The following results follow from Theorem 4.2 in [25]:

Theorem 5.1. *Let $Q_1 \leq Q_2 \leq \dots$ be a sequence of closed semibounded from below sesquilinear forms on \mathcal{H} . Then*

$$Q_\infty(f_1, f_2) := \lim_{n \rightarrow \infty} Q_n(f_1, f_2)$$

with

$$\text{Dom}(Q_\infty) := \left\{ f \in \bigcap_{n \in \mathbb{N}} \text{Dom}(Q_n) : \sup_{n \in \mathbb{N}} Q_n(f, f) < \infty \right\}$$

is a closed semibounded from below sesquilinear form Q_∞ in \mathcal{H} , and one has

$$e^{-tS_{Q_n}} P_{Q_n} \rightarrow e^{-tS_{Q_\infty}} P_{Q_\infty} \quad \text{strongly in } \mathcal{L}(\mathcal{H}) \text{ as } n \rightarrow \infty, \text{ for all } t \geq 0.$$

Theorem 5.2. *Let $Q_1 \geq Q_2 \geq \dots$ be a sequence of closed semibounded sesquilinear forms on \mathcal{H} . Define a nonnegative sesquilinear form in \mathcal{H} given by*

$$Q'(f_1, f_2) := \lim_{n \rightarrow \infty} Q_n(f_1, f_2), \quad \text{Dom}(Q') := \bigcup_{n \in \mathbb{N}} \text{Dom}(Q_n)$$

and let Q_∞ denote the form on \mathcal{H} given by the closure of $\text{reg}(Q')$ in \mathcal{H} . Then one has

$$e^{-tS_{Q_n}} P_{Q_n} \rightarrow e^{-tS_{Q_\infty}} P_{Q_\infty} \quad \text{strongly in } \mathcal{L}(\mathcal{H}) \text{ as } n \rightarrow \infty, \text{ for all } t \geq 0.$$

Remark 5.3. Note that in the above situation Q_∞ is the closure of Q' if Q' is closable, and $Q_\infty = Q'$ if Q' is even closed.

References

- [1] T. Aubin. Espaces de Sobolev sur les varietes riemanniennes. Bull. Sci. Math. (2) 100 (1976), no. 2, 149–173.
- [2] B. Booß-Bavnbek, K. P. Wojciechowski. Elliptic boundary problems for Dirac operators. Mathematics: Theory & Applications. Birkhäuser Boston, Inc., Boston, MA, 1993.
- [3] B. K. Driver, A. Thalmaier. Heat equation derivative formulas for vector bundles. J. Funct. Anal. 183 (2001), no. 1, 42–108.
- [4] M. Fukushima, Y. Oshima, M. Takeda. Dirichlet forms and symmetric Markov processes. De Gruyter Studies in Mathematics, 19. Walter de Gruyter & Co., Berlin, 1994.
- [5] W. G. Faris. Self-adjoint operators. Lecture Notes in Mathematics, Vol. 433. Springer-Verlag, Berlin-New York, 1975.
- [6] A. Grigoryan. Heat kernels on weighted manifolds and applications. The ubiquitous heat kernel. Contemp. Math., 398, 93–191, Amer. Math. Soc., Providence, RI, 2006.

- [7] A. Grigoryan. Heat kernel and analysis on manifolds. AMS/IP Studies in Advanced Mathematics, 47. American Mathematical Society, Providence, RI; International Press, Boston, MA, 2009.
- [8] B. Güneysu. Heat kernels in the context of Kato potentials on arbitrary manifolds. *Potential Anal.* 46 (2017), no. 1, 119–134.
- [9] B. Güneysu. On generalized Schrödinger semigroups. *J. Funct. Anal.* 262 (2012), no. 11, 4639–4674.
- [10] B. Güneysu. Nonrelativistic hydrogen type stability problems on nonparabolic 3-manifolds. *Ann. Henri Poincaré* 13 (2012), no. 7, 1557–1573.
- [11] B. Güneysu. Covariant Schrödinger semigroups on Riemannian manifolds. *Operator Theory: Advances and Applications*, 264. Birkhäuser/Springer, Cham, 2017. xviii+239 pp.
- [12] I. L. Hedberg. Spectral synthesis and stability in Sobolev spaces. *Euclidean harmonic analysis (Proc. Sem., Univ. Maryland, College Park, Md., 1979)*, pp. 73–103, *Lecture Notes in Math.*, 779, Springer, Berlin, 1980.
- [13] I. W. Herbst, Z. X. Zhao. Sobolev spaces, Kac-regularity, and the Feynman-Kac formula. *Seminar on Stochastic Processes, 1987 (Princeton, NJ, 1987)*, 171–191, *Progr. Probab. Statist.*, 15, Birkhäuser Boston, Boston, MA, 1988.
- [14] R. Grummt, M. Kolb. Essential selfadjointness of singular magnetic Schrödinger operators on Riemannian manifolds. *J. Math. Anal. Appl.* 388 (2012), no. 1, 480–489.
- [15] D. Guidetti, B. Güneysu, D. Pallara. L^1 -elliptic regularity and $H = W$ on the whole L^p -scale on arbitrary manifolds. *Ann. Acad. Sci. Fenn. Math.* 42 (2017), no. 1, 497–521.
- [16] V. Guillemin, D. Pollack. *Differential topology*. Prentice-Hall, Inc., Englewood Cliffs, N.J., 1974.
- [17] M. W. Hirsch. *Differential topology*. Corrected reprint of the 1976 original. *Graduate Texts in Mathematics*, 33. Springer-Verlag, New York, 1994
- [18] E. P. Hsu. *Stochastic analysis on manifolds*. *Graduate Studies in Mathematics*, 38. American Mathematical Society, Providence, RI, 2002.
- [19] K. Kuwae, M. Takahashi. Kato class functions of Markov processes under ultracontractivity. *Potential theory in Matsue, 193–202*, *Adv. Stud. Pure Math.*, 44, Math. Soc. Japan, Tokyo, 2006.
- [20] K. Kuwae, M. Takahashi. Kato class measures of symmetric Markov processes under heat kernel estimates. *J. Funct. Anal.* 250 (2007), no. 1, 86–113.
- [21] H.B. Jr. Lawson, M.-L. Michelsohn. *Spin geometry*. *Princeton Mathematical Series*, 38. Princeton University Press, Princeton, NJ, 1989.
- [22] M. Lesch. *Operators of Fuchs type, conical singularities, and asymptotic methods*. *Teubner-Texte zur Mathematik [Teubner Texts in Mathematics]*, 136. B. G. Teubner Verlagsgesellschaft mbH, Stuttgart, 1997.
- [23] L.I. Nicolaescu. *Lectures on the geometry of manifolds*. Second edition. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007.
- [24] L. F. Pessoa, S. Pigola, A. G. Setti. Dirichlet parabolicity and L^1 -Liouville property under localized geometric conditions. <https://arxiv.org/abs/1607.06483>
- [25] B. Simon. A canonical decomposition for quadratic forms with applications to monotone convergence theorems. *J. Funct. Anal.* 28 (1978), no. 3, 377–385.
- [26] P. Stollmann. *Formtechniken bei Schrödingeroperatoren*. Diploma thesis, Munich 1985.
- [27] P. Stollmann, J. Voigt. Perturbation of Dirichlet forms by measures. *Potential Anal.* 5 (1996), no. 2, 109–138.
- [28] D. Stroock. The Kac approach to potential theory: Part I. *J. Mathematics and Mechanics*. vol. 16. no. 8. 829–852 (1967).
- [29] M. E. Taylor. *Partial differential equations. I. Basic theory*. *Applied Mathematical Sciences*, 115. Springer-Verlag, New York, 1996